



# A Role for Central Spindle Proteins in Cilia Structure and Function

Katherine R. Smith,<sup>1</sup> Esther K. Kieserman,<sup>1</sup> Peggy I. Wang,<sup>2,3</sup> Sander G. Basten,<sup>4</sup>  
Rachel H. Giles,<sup>4</sup> Edward M. Marcotte,<sup>3,5,6</sup> and John B. Wallingford<sup>1,6,7\*</sup>

<sup>1</sup>Section of Molecular Cell and Developmental Biology, University of Texas at Austin, Austin, Texas

<sup>2</sup>Department of Biomedical Engineering, University of Texas at Austin, Austin, Texas

<sup>3</sup>Center for Systems and Synthetic Biology, University of Texas at Austin, Austin, Texas

<sup>4</sup>Department Medical Oncology, University Medical Center Utrecht, Utrecht, The Netherlands

<sup>5</sup>Department of Chemistry and Biochemistry, University of Texas at Austin, Austin, Texas

<sup>6</sup>Institute for Cellular and Molecular Biology, University of Texas at Austin, Austin, Texas

<sup>7</sup>Howard Hughes Medical Institute, University of Texas at Austin, Austin, Texas

Received 12 November 2010; Accepted 28 November 2010

Monitoring Editor: Bruce Goode

**Cytokinesis and ciliogenesis are fundamental cellular processes that require strict coordination of microtubule organization and directed membrane trafficking. These processes have been intensely studied, but there has been little indication that regulatory machinery might be extensively shared between them. Here, we show that several central spindle/midbody proteins (PRC1, MKLP-1, INCENP, centriolin) also localize in specific patterns at the basal body complex in vertebrate ciliated epithelial cells. Moreover, bioinformatic comparisons of midbody and cilia proteomes reveal a highly significant degree of overlap. Finally, we used temperature-sensitive alleles of *PRC1/spd-1* and *MKLP-1/zen-4* in *C. elegans* to assess ciliary functions while bypassing these proteins' early role in cell division. These mutants displayed defects in both cilia function and cilia morphology. Together, these data suggest the conserved reuse of a surprisingly large number of proteins in the cytokinetic apparatus and in cilia.** © 2010 Wiley-Liss, Inc.

**Key Words:** ciliogenesis, cytokinesis, PRC1, INCENP, MKLP-1, bioinformatics, cilia midbody

## Introduction

Cilia are microtubule-based organelles that project from cells, and mutations of genes associated with ciliogenesis underlie a broad spectrum of disorders in

humans [Marshall, 2008; Sharma et al., 2008]. As such, understanding the full complement of proteins required for establishing and maintaining cilia structure and function will be important. Genomic and proteomic approaches have defined the basal body and cilia proteomes, but many proteins essential for ciliogenesis are not present in these datasets [Gherman et al., 2006; Inglis et al., 2006; Park et al., 2006; Hayes et al., 2007; Gray et al., 2009]. Functional screens might be expected to reveal these other important ciliogenesis factors [e.g., Kim et al., 2010], but many proteins have pleiotropic functions that may render their role in cilia invisible to such screens. Indeed, recent reports demonstrate exocytic roles for some proteins required for ciliogenesis [Finetti et al., 2009; Gray et al., 2009].

Cilia assembly and function require strictly coordinated microtubule organization and membrane delivery [Sorokin, 1968; Nachury et al., 2007; Park et al., 2008; Gray et al., 2009]. Cells face a similar challenge during cytokinesis, as proteins in the central spindle must both organize the microtubule bundles and direct membrane traffic into the nascent cleavage furrow [Gromley et al., 2005; Albertson et al., 2008; Glotzer, 2009]. In fact, a small number of regulatory proteins have been implicated in both processes (e.g. CP110, the exocyst, BBS6, RhoA; [Gromley et al., 2005; Kim et al., 2005; Pan et al., 2007; Spektor et al., 2007; Park et al., 2008; Shah et al., 2008; Zuo et al., 2009]).


The protein network that includes PRC1, INCENP, MKLP-1, and centriolin acts at the central spindle and midbody and has been studied extensively for its role in cytokinesis [Jiang et al., 1998; Kurasawa et al., 2004; Gromley et al., 2005; Kieserman et al., 2008; Glotzer, 2009]. In this paper, we report novel roles for these proteins, showing that they localize to basal bodies in ciliated epithelial cells and control cilia function and morphology.

Additional Supporting Information may be found in the online version of this article.

Esther K. Kieserman's present address is Department of Molecular and Cell Biology, University of California, Berkeley, California.

\*Address correspondence to: John B. Wallingford, 1 University Station C1000, Austin, TX 78712, USA. E-mail: wallingford@mail.utexas.edu

Published online 7 December 2010 in Wiley Online Library (wileyonlinelibrary.com).



**Fig. 1. Central spindle/midbody proteins localize to basal bodies in multiciliated epithelial cells.** (A) SEM of the ciliated epidermis of the *Xenopus laevis* tadpole showing a multiciliated cell and surrounding mucus secreting goblet cells. (B) Merged image of immunostaining of PRC1 (green) and  $\alpha$ -tubulin (red) in the epidermis of a *X. laevis* tadpole. Arrow points to a ciliated cell with cilia protruding from the surface and a punctate pattern of PRC1 under the cilia. Arrowhead points to a dividing cell in mid-telophase with PRC1 localization at the central spindle. (b') Same image as B showing only PRC1 localization. (b'') Same image as B showing only  $\alpha$ -tubulin localization. (C) Immunostaining of MKLP-1 in a ciliated cell. (c') Merged image of immunostaining of MKLP-1 and  $\alpha$ -tubulin to reveal cilia. (D) Immunostaining of INCENP in a ciliated cell. (d') Merged image of immunostaining of INCENP and  $\alpha$ -tubulin to reveal cilia. (E, e') Immunostaining of PRC1 (green) and acetylated tubulin (red) in human RPE serum starved cells shows PRC1 localizing to the base of primary cilia. (F) Immunostaining of MKLP-1 (green) and acetylated tubulin (red) in RPE cells also shows MKLP-1 localizing to the base of primary cilia. Nuclei stained with Hoechst. Scale bars B = 10  $\mu$ m, C, D = 3  $\mu$ m, E = 5  $\mu$ m, and e', F = 2  $\mu$ m.


Our data suggest the possibility that a surprising amount of cytokinetic regulatory machinery may also serve important functions related to cilia.

## Results and Discussion

We have developed a method to identify novel ciliogenesis factors based upon differential gene expression in the multiciliated epithelial cells of the *Xenopus* embryo epidermis (Fig. 1A; see also Hayes et al. [2007]). Extending this system to protein localization, we identified the microtubule-bundling protein PRC1 as localizing prominently to the

base of cilia in multiciliated cells (Figs. 1B and 1b' arrow). PRC1 is one member of a well-defined protein network acting at the central spindle and the midbody during cytokinesis [Glotzer, 2005; Glotzer, 2009; Jiang et al., 1998] (Fig. 1B, arrowhead). These proteins organize microtubule bundles, position the cleavage furrow, and also trigger vesicle recruitment for cell abscission [Jiang et al., 1998; Glotzer, 2005; Gromley et al., 2005; Glotzer, 2009].

The localization of PRC1 near basal bodies was surprising, since central spindle proteins have not previously been implicated in ciliogenesis. We therefore asked if




**Fig. 2. Spatially distinct localization of central spindle/midbody proteins at the basal body.** (A) Early tailbud stage *Xenopus laevis* embryo and diagram showing the planar polarization of cilium structures in the multiciliated cells of the epithelium, with the rootlet (green) positioned on the anterodorsal face of the basal body (red). (B) PRC1 (green) and  $\gamma$ -tubulin (red) immunostaining shows that PRC1 is proximal to the basal body on the anterior dorsal side, (b')  $x$ - $y$  view of PRC1 and  $\gamma$ -tubulin with diagram below showing section views. (b'')  $x$ - $z$  plane view. (C) Coimmunostaining against MKLP-1 (green) and  $\gamma$ -tubulin (red) shows MKLP-1 proximal to but not colocalizing with the basal body on the anterior dorsal side. (c')  $x$ - $y$  plane view of MKLP-1 and  $\gamma$ -tubulin localization with diagram below showing section views. (c'')  $x$ - $z$  plane view. (D) MKLP-1 colocalizes with rootlet marker CLAMP in a polarized manner. (d')  $x$ - $y$  view showing colocalization between MKLP-1 and CLAMP, with diagram below showing  $x$ - $z$  and  $x$ - $y$  slices and overlap. (d'')  $x$ - $z$  plane view. Scale bars B-D = 5  $\mu$ m and b'-d'' = 1  $\mu$ m.

other proteins in this network also localized to basal bodies. We found that MKLP-1, INCENP, centriolin, CENP-E, and Survivin also localized strongly to the base of cilia in *Xenopus* multiciliated cells (Figs. 1C, 1C', 1D, 1d', 3C, and 3c''; data not shown).

The PRC1 and MKLP1 antibodies used for immunostaining in *Xenopus* were ineffective in western blots with *Xenopus* embryo lysates, and some antibodies made in rabbits can nonspecifically label centrosomes. Therefore, to verify the observed localization patterns, we obtained





**Fig. 3. INCENP colocalizes with the rootlet and Centriolin with the basal body.** (A) Immunostaining of INCENP (green). (a') Same image as A showing immunostaining of  $\gamma$ -tubulin (red). (a'') Merged image INCENP and  $\gamma$ -tubulin. (a''') Immunostaining shows INCENP localizing proximal to, but not with,  $\gamma$ -tubulin. (B) Immunostaining of INCENP (green). (b') Same image as B showing CLAMP-RFP localization. (b'') Merged image of INCENP and CLAMP-RFP. (b''') Immunostaining shows INCENP colocalizing with CLAMP-RFP. (C) Centriolin (green) localization in a multiciliated cell. (c') Same image as C showing  $\gamma$ -tubulin (red) localization. (c'') Merged image of Centriolin and  $\gamma$ -tubulin shows colocalization. Scale bars a'–c' = 5  $\mu$ m and a'', b'' = 1  $\mu$ m.

additional antibodies against these proteins (see Materials and Methods) and tested them in a different cell type, human RPE cells. Immunostaining for PRC1, MKLP-1, INCENP, and Survivin revealed that each of these proteins prominently localized to the base of primary cilia in serum-starved human RPE cells in culture (Figs. 1E, 1e', and 1F; Supporting Information Fig. 1, and data not shown). These data therefore suggest that central spindle proteins localize to the base of both specialized motile

cilia in *Xenopus* multiciliated cells and to the base of primary cilia in human RPE cells.

We further defined the localization of these central spindle proteins in multiciliated cells by coimmunostaining with  $\gamma$ -tubulin, a marker of the basal body. Interestingly, PRC1 did not localize to the basal body itself, but rather lay immediately adjacent to it in a planar polarized fashion (Figs. 2B, 2b', and 2b''). This pattern was intriguing because planar polarization of basal bodies in multiciliated

cells is central to directed ciliary beating and fluid flow (Fig. 2A) [Frisch and Farbman, 1968; Mitchell et al., 2007; Park et al., 2008].

Basal body planar polarity is manifested in the stereotyped positioning of accessory structures, such as the ciliary rootlet and basal foot [Frisch and Farbman, 1968; Mitchell et al., 2007]. In *Xenopus* multiciliated cells, rootlets are consistently positioned anterodorsally to the basal body, opposite from the direction of fluid flow (Fig. 2A; [Mitchell et al., 2007; Park et al., 2008]). In X/Y projections, PRC1, MKLP-1, and INCENP were consistently present on the anterodorsal side of the  $\gamma$ -tubulin-positive basal body (Figs. 2B, 2b', 2C, 2c', 3A, and 3a''), suggesting that these proteins may localize in the vicinity of the ciliary rootlet.

Rootlets in *Xenopus* multiciliated cells project basolaterally into the cytoplasm [Steinman, 1968]. Likewise, X/Z projections from 3D datasets revealed that PRC1 and MKLP1 signals extended basolaterally into the cytoplasm from the anterodorsal face of the  $\gamma$ -tubulin-positive basal body (Figs. 2b'' and 2c''). Moreover, the microtubule-binding protein, CLAMP marks the vicinity of the ciliary rootlet [Park et al., 2008], and coimmunostaining demonstrated that MKLP-1 and INCENP each partially colocalized with CLAMP-RFP (Figs. 2D, 2d'', 3B, and 3b''). Together, these data suggest that central spindle proteins localize in a planar polarized fashion at the base of cilia in the vicinity of the rootlet.

The rootlet is implicated as a key site where vesicular traffic is coordinated with microtubule-based structures during ciliogenesis [Fariss et al., 1997; Yang and Li, 2005; Park et al., 2008; Gray et al., 2009]. The localization of central spindle proteins near this site suggested a potential parallel between ciliogenesis and cytokinesis, as the central spindle also governs vesicle delivery during cytokinesis [Gromley et al., 2005; Albertson et al., 2008]. To further explore this parallel, we examined centriolin, which localizes to the midbody ring, immediately adjacent to the central spindle proteins, and which directs the recruitment of vesicles [Gromley et al., 2005]. We found that centriolin was also localized to the base of cilia in multiciliated cells (Figs. 3C and 3c''). Interestingly, centriolin did not localize near the rootlet, but rather colocalized with  $\gamma$ -tubulin at the basal body itself (Fig. 3c''). In summary, localization experiments in *Xenopus* multiciliated cells revealed that several components of the protein network governing cytokinesis localize to spatially distinct domains at the basal body that reflect to some degree their function in the midbody.

To more comprehensively explore the potential link between cytokinesis and ciliogenesis, we used bioinformatic approaches to compare the midbody proteome, as defined by proteins known to localize to the midbody [Skop et al., 2004; Ren et al., 2009] with the cilia/basal body proteome, as defined by mass spectrometry and comparative genomics


[Gherman et al., 2006; Inglis et al., 2006]. We found that the two proteomes overlap significantly (Fig. 4A, blue overlap;  $P \leq 10^{-14}$ , hypergeometric probability). The cilia/basal body proteome includes proteins identified directly, as well as many predicted by comparative genomics and other means [Gherman et al., 2006; Inglis et al., 2006]. To more stringently examine the overlap of these proteomes, we re-examined the overlap using only centrosome or basal body proteins as determined directly by mass spectrometry. Again, we observed far greater overlap than could be expected by chance (Fig. 4A, green overlap,  $P \leq 10^{-13}$ , hypergeometric probability).

To ask if this overlap between midbody and basal body proteomes is biologically relevant, we examined the positioning of these proteins in a functional gene network. This network is composed of functional interactions among human genes (determined as a probabilistic combination of evidence for coregulation, physical and/or genetic interactions, coevolution, and interactions between orthologs in other species) and serves as a useful indicator of proteins operating in the same biological processes and pathways [Lee et al., 2008b; Gray et al., 2009; Li et al., 2009]. Proteins found in the midbody and cilia/basal body proteomes were much more strongly interconnected within the probabilistic network than would be expected by chance (Fig. 4B;  $P \leq 10^{-8}$ , Z-score test). In fact, 80% of the overlapping proteins clustered together tightly within the network (Fig. 4B).

This cluster of proteins included not only the few already known to play roles in both cytokinesis and ciliogenesis [e.g. exocyst, RhoA [Glutzer, 2005; Gromley et al., 2005; Pan et al., 2007; Park et al., 2008; Zuo et al., 2009] (Fig. 4B, red/brown)], but also several additional proteins of interest. For example, Plk1 was present in this cluster within the network (Fig. 4B, purple); Plk1 interacts with PRC1 at the central spindle [Neef et al., 2007] but has not been previously implicated in cilia. The cluster also contains several MAP kinases, whose connection in this network may shed light on the mechanisms by which FGF/MAP kinase signaling, which is well-known to control cell division, also controls cilia length [Berman et al., 2003; Burghoorn et al., 2007; Neugebauer et al., 2009] (Fig. 4B, yellow). Finally, the cluster contains a known human ciliopathy gene, *Dynl1* [Horvath et al., 2005].

PRC1 and other central spindle proteins are not present in this cluster because they have not previously been associated with cilia or basal bodies. In order to link our bioinformatic analyses to our immunostaining results, we added PRC1 and MKLP-1 to the overlapping proteome set and re-examined the connectivity in our network. Both PRC1 and MKLP-1 were tightly embedded in this cluster of connected genes (Fig. 5).

Together with our immunostaining data, these bioinformatic analyses suggest the possibility that a functionally analogous protein network acts at both the central spindle/




**Fig. 4. Bioinformatic analyses reveal a large set of common proteins involved in cytokinesis and ciliogenesis.** (A) Venn diagram showing the overlap between the midbody and basal body/ciliary proteomes. The overlap between midbody proteome and either core and extended basal body/cilia proteomes (see Supporting Information) is significantly greater than expected by chance (hypergeometric test). (B) A network diagram of gene-gene associations is drawn for proteins present in both the midbody proteome and core (green squares) or extended (blue circles) basal body/cilia proteome. Lines connect proteins likely to function together on the basis of coexpression, coregulation, and/or protein-protein interactions, as described in the Supporting Information. The core cilia proteome is preferentially connected by network associations to midbody proteins ( $P \leq 10^{-8}$ , see Supporting Information). This cluster contains several notable proteins, including exocyst subunits (red), RhoA (brown), several MAP kinases (yellow), PLK1 (purple), and the human ciliopathy gene *DynLL1* (gray).

midbody to control cytokinesis and at the basal body to control cilia structure or function. However, the essential role for these proteins in cell division would render their role in cilia invisible to standard loss-of-function approaches *in vivo*. To circumvent this problem, we obtained a temperature-sensitive mutant of PRC1 in the worm, *C. elegans*. This *PRC1* allele, *spd-1*, has normal function at 15°C but is nonfunctional at 25°C [O’Connell et al., 1998; Verbrugge and White, 2004]. Worms carrying this mutation were reared at the permissive temperature through early develop-

ment and then shifted to the restrictive temperature at the time when cilia are thought to begin assembling on sensory neurons [Perkins et al., 1986; Swoboda et al., 2000]. We then assessed the function and morphology of neuronal cilia in these animals.

The proper functioning of neuronal sensory cilia is essential for worms to detect chemoattractants [Perkins et al., 1986], so we initially assayed the role of PRC1 in cilia function by scoring chemotaxis of *spd-1* worms at permissive and restrictive temperatures. Because *spd-1* mutant worms






**Fig. 5. PRC1 and MKLP-1 are tightly linked in the cluster of cytokinesis and ciliogenesis genes in a probabilistic gene network.** A network diagram of gene-gene associations is drawn for PRC1, MKLP-1 and the proteins present in both the midbody proteome and core (green squares) or extended (blue circles) basal body/cilia proteome. Lines connect proteins likely to function together on the basis of coexpression, coregulation, and/or protein-protein interactions, as described in the Supporting Information.

display a mild uncoordinated phenotype [O’Connell et al., 1998], we designed a quantification method that uncouples effects on movement and chemotaxis (Fig. 6A). Worms were placed equidistant from circles containing the attractant or containing no attractant (Fig. 6A), and chemotaxis was quantified as the number of worms successfully moving into the chemoattractant circle as compared to the number successfully moving into the empty circle. In this assay, *spd-1* worms at the permissive temperature chemotaxed normally (Fig. 6B). Importantly however, when shifted to the restrictive temperature, *spd-1* worms failed to chemotax (Fig. 6B). The chemotaxis defect observed with *spd-1* mutant worms at the restrictive temperature was comparable to that of *daf-19* depleted worms, which are known to lack cilia (Fig. 6B) [Perkins et al., 1986; Swoboda et al., 2000].

To confirm a defect in cilia function, we next exploited the finding that functional cilia mediate the uptake of dye

by sensory neurons (Fig. 6C) [Hedgecock et al., 1985]. We observed that many neuronal cell bodies in *spd-1* worms at restrictive temperatures failed to fill with dye (Fig. 6D, arrow). Moreover, those neurons in *spd-1* mutants that did fill were consistently less brightly labeled than were control neurons (Fig. 6D, arrowhead). We quantified dye intensity in several neurons from several animals and observed that *spd-1* mutant worms at restrictive temperature displayed a 43% reduction in fluorescence intensity as compared to controls (Supporting Information Fig. 2B). Similar phenotypes - both loss of dye filling and reduction in intensity of dye filling - are observed in *bona fide* cilia mutants in *C. elegans* [Perkins et al., 1986]. Thus, two distinct assays indicate that cilia function was impaired in the absence of wild-type PRC1.

To examine the role of PRC1 in governing cilia morphology, we crossed the *spd-1* worms into the *str-1::GFP*



**Fig. 6. PRC1 is essential for cilia function in *C. elegans*.** (A) Diagram on left shows the experimental set-up: worms were placed in the middle of the plate and then the number present in a circle with chemoattractant (green) or without (blue) was scored. (B) Graph shows that *spd-1* mutants at 25° chemotax ineffectively compared to *spd-1* mutants at 15°C ( $P < 0.0001$ , Fisher's exact test) and MT8189 controls ( $P < 0.0001$ , Fisher's exact test), and this defect is comparable to worms fed RNAi against *daf-19* ( $P$ -value n.s.), which lack cilia [Perkins et al., 1986]. For the chemotaxis assay  $n > 200$  for all strains. (C) FITC dye filling in neurons of a control worm. (D) Defective neuronal filling with FITC dye in *spd-1* worms at 25°C. Outlines in C and D represent approximate position of neuronal cell bodies. Some cells fail to fill (arrow), while others display reduced filling (arrowhead). Overall, fluorescence intensity in *spd-1* mutant worms at 25°C was reduced to 57% of control levels (see Supporting Information Fig. 4).

strain, with which the cilia on AWB neurons can be visualized [Mukhopadhyay et al., 2007, 2008]. The AWB cilia develop a complex architecture that is perturbed by mutation of genes involved in ciliogenesis, thus providing a sensitive assay for candidate genes [Mukhopadhyay et al., 2007, 2008; Murayama et al., 2005]. AWB cilia typically display a two-pronged structure, though there is some variability in control worms, and cilia with ectopic branches are occasionally observed (Figs. 7A, 7B, 7E, and 7I; and see Mukhopadhyay et al., [2007, 2008]). In *spd-1(str-1::GFP)* mutant worms at restrictive temperature, we observed a dramatic increase in the number of cilia with ectopic branches (Figs. 7C, 7E, 7G, and 7I). Moreover, we observed an increased frequency of large bulges on sensory cilia of mutant worms (Figs. 7H and 7I). These phenotypes in AWB cilia, both bulges and ectopic projections, have been previously observed following mutation of genes involved in ciliogenesis [Mukhopadhyay et al., 2007, 2008].

Finally, because PRC1 acts at the central spindle together with MKLP-1 [Kurasawa et al., 2004] and because both proteins localize to the base of cilia (Figs. 1B and 1C), we assessed the role of MKLP-1 in AWB cilia. Using the temperature-sensitive MKLP-1 mutant *zen-4*, we


found that loss of MKLP-1 resulted in a significantly increased number of ectopically branched AWB cilia (Figs. 7D and 7I), consistent with our results for PRC1/*spd-1*.

## Conclusions

Our data demonstrate that PRC1 and MKLP-1 are required for normal AWB cilia morphology in *C. elegans*. Coupled to the spatially distinct patterns of localization of PRC1, MKLP-1, INCENP, centriolin, and other central spindle/midbody proteins at basal bodies in *Xenopus* motile cilia and in human primary cilia, our data suggest that these proteins play a conserved role in regulating cilia structure and function.

The function of the central spindle proteins in relation to cilia remains unknown. One role for these proteins in the central spindle is microtubule bundling, and so it is possible that they are required for centrosome or basal body structure; indeed, MKLP-1 and PRC1 can localize to centrosomes. However, in postmitotic multiciliated cells in *Xenopus*, we found these proteins to localize adjacent to, but not in, the basal body, suggesting that they may serve a different function. Another possibility is that these proteins act in ciliary disassembly, as the Fa2p kinase





**Fig. 7. PRC1 and MKLP1 control morphology of sensory cilia in *C. elegans*.** (A,B) Wild-type worms expressing GFP behind the *str-1* promoter in the AWB neurons. The majority (81%) of wild-type *str-1::GFP* worms have no ectopic projections (blue, E) on their AWB cilia. In *spd-1* worms at 25°C (C), however, over half (56%) have small (F) to large (G) ectopic projections (green, purple, asterisk), while a minority have an additional phenotype of GFP accumulation at the distal tip (H, orange). In *zen-4* worms (D) at 25°C there is also an increase in ectopic projections (42%). (I) Percentages of worms with cilia morphology. For cilia morphology *str-1::GFP*,  $n = 27$ , *spd-1(str-1::GFP)*,  $n = 43$ , *zen-4(str-1::GFP)*,  $n = 38$ . For A–C scale bars = 5  $\mu\text{m}$ , D–G scale bars = 2  $\mu\text{m}$ .

regulates ciliary disassembly in *Chlamydomonas* but localizes to midbodies when expressed in mammalian cells [Mahjoub et al., 2004].

Another possible role for the central spindle proteins is in recruitment of exocyst-mediated vesicle traffic to the cytokinetic furrow [Gromley et al., 2005]. Given the localization of exocyst components to the base of cilia and their implication in ciliogenesis [Park et al., 2008; Zuo et al., 2009], it is tempting to suggest that the central spindle proteins may influence cilia-related vesicle traffic. This model would be consistent with our localization data, as the rootlet is implicated as a site of vesicle trafficking [Fariss et al., 1997; Yang and Li, 2005; Park et al., 2008; Gray et al., 2009]. While the mechanisms of action must be explored further, the identification of this new connection between the cytokinetic apparatus and cilia should accelerate our understanding of both, as further study in one context may now provide important new insights for understanding the other (Supporting Information Fig. 3).

Finally, our data may shed light on the etiology of human ciliopathies such as Bardet-Biedel Syndrome, as the BBS6 protein localizes prominently to the midbody during cytokinesis [Kim et al., 2005]. Indeed, while mutation of this gene in humans leads to a disease that is classified as a ciliopathy, experimental manipulation of BBS6 has been found to have relatively modest effects on cilia

morphology and function [Shah et al., 2008]. By contrast, knockdown of BBS6 prevents cell abscission during cytokinesis [Kim et al., 2005]. Likewise, knockdown of the intraflagellar transport protein IFT27 leads to both flagellar defects and cell division defects [Qin et al., 2007], and this protein has been reported to localize to vesicles associated with the cytokinetic furrow [Baldari and Rosenbaum, 2010]. Together with these findings, our data raise the possibility of a connection between cytokinesis defects and ciliopathy phenotypes. Thus, while the pleiotropic roles of central spindle proteins will make the study of their roles in cilia challenging, further exploration of the functional connection between the cytokinetic machinery and cilia should be exciting.

## Material and Methods

### *Xenopus* Embryo Isolation and Injection

Female *Xenopus laevis* were injected with 700 ml of human chorionic gonadotropin hormone and stored at 18°C overnight. The next day, oocytes were isolated from mothers and fertilized [Sive et al., 2000]. Embryos were dejellied with a 3% cysteine solution in 1/3X MMR. To image CLAMP localization embryos were injected on the ventral side of the embryo at the four-cell stage with 300 pg/per injection of CLAMP-RFP

mRNA made using the mMessage kit (Ambion). Embryos were grown to stage 13, early neurula, and stage 30, tailbud, before fixation.

### Immunohistochemistry on *Xenopus* Embryos

Immunostaining for central spindle and passenger proteins was performed essential as previously described [Lee et al., 2008a].

Embryos were grown to appropriate stages and fixed in Dent's Fix (80% methanol/20% DMSO) [Becker and Gard, 2006] overnight at 4°C. Embryos were fully dehydrated in methanol and stored at -20°C overnight. Embryos were slowly rehydrated by 5 min washes in 75% methanol in PTW (phosphate-buffered saline [PBS] + 1% Tween-20), 50% methanol in PTW, 25% methanol in PTW, and finally PTW. Embryos were bleached in a light box in a solution of 10% H<sub>2</sub>O<sub>2</sub>, 0.5% formamide, and 2.5% 2× SCC. Embryos were then washed 3× 5 min in Tris-buffered saline + 1% Tween-20 (TBST). Embryos were blocked in 300 ml of TBS + 10% FBS + 5% DMSO. Primary antibody was added; rabbit-anti-PRC1 (1:20; BioLegend), rabbit-anti-MKLP1 (1:200) [Potapova et al., 2006], rabbit-anti-INCENP (1:800) [Potapova et al., 2006], Aurora-B (1:50) [Potapova et al., 2006], rabbit-anti-centriolin serum (1:100) [Gromley et al., 2005], or mouse-anti-γ-tubulin (1:250; Abcam ab27076). Embryos were incubated overnight at 4°C. Embryos were washed 5× 1 hr in TBST at RT the next day. Embryos were then blocked as before for 1 hr and anti-rabbit Alexa fluor 488 (1:250) and anti-mouse Alexa fluor 555 (1:250) were added. Embryos were incubated again over night at 4°C and washed 5× 1 hr in TBST at RT the next day. For embryos that were being imaged for α-tubulin localization TBST was removed and 300 ml of mouse-anti-α-tubulin antibody + blocking solution was added (1:250). Embryos were incubated for 1 hr at room temperature. Embryos were washed again for 3× 1 hr washes in TBST. Secondary antibodies were added as above. In some cases during the last wash a dilution of (1:2000) of DAPI was added to label nuclei.

### Image Acquisition of *Xenopus* Embryos

All imaging was performed using a Zeiss LSM5 PASCAL microscope with either 40x planNeofluar (1.3na), 63x planApochromat (1.4na) oil immersion objectives or 100x Fluor (1.3na). Confocal optical slices were collected and maximum intensity projections made with Zeiss LSM5 software. For presentation purposes, some images have been processed with Adobe Photoshop.

### Cell Culture and Imaging

RPE-hTERT cells were cultured in DMEM/F12 (10% FBS/PS/Glut). Confluent cells were ciliated by serum starvation in (0.2% FCS) for 48 hr. Cells were fixed in

methanol (ice cold) for 5 min on ice or 4% PFA for 15 min at room temperature. Following fixation cells were blocked for 1 hr in 3% BSA/0.1% Triton-X100 and incubated with primary antibodies; rabbit-anti-PRC1 (1:50; Abcam, ab21437), rabbit-anti-MKLP-1 (1:100; Santa Cruz, sc-22793; H-110), and anti-acetylated-tubulin (1:10,000; Sigma-Aldrich T6793; clone 6-11B-1) 2–4 hr at RT in 3% BSA. Secondary antibody incubation was performed for 1 hr at room temperature using Jackson ImmunoResearch secondary antibodies (1:500) and Hoechst (1:100,000; 33342, Invitrogen). Samples were embedded in Fluoromount-G (Beckman Coulter). Imaging was performed using an Everest deconvolution workstation (Intelligent Imaging Innovations) equipped with a Zeiss AxioImager.Z1 microscope and a CoolSnapHQ-cooled CCD camera (Roper Scientific) and a 63x PlanApochromat, NA 1.4 objective and processed using SlideBook 5 (Intelligent Imaging Innovations) and ImageJ.

### Proteome Sets

Cilia proteins were downloaded from the ciliary proteome database at <http://v3.ciliaproteome.org/cgi-bin/index.php> [Gherman et al., 2006]. The core basal body proteome set was defined as centrosome and centriole proteins identified by mass spectrometry (MS) [Andersen et al., 2003; Keller et al., 2005]. For the extended set, basal body, flagellum, and cilium proteins identified by either MS or comparative genomics were also included [Avidor-Reiss et al., 2004; Li et al., 2004; Liu et al., 2007]. Human midbody proteins were downloaded from MiCroKit 3.0, a manually curated database of protein subcellular localizations, experimentally verified by fluorescence microscopy (<http://microkit.biocuckoo.org>, [Ren et al., 2009]). The set of all human Entrez proteins were retrieved from the Ensembl database at [www.ensembl.org](http://www.ensembl.org) [Hubbard et al., 2009].

### Proteome Overlap Analysis

The hypergeometric distribution was used to assess if the midbody and cilia proteomes share a significant number of proteins. The cumulative *P*-value was calculated using

$$p_{cdf} = \sum_{i=k}^{\min\{m,n\}} \frac{\binom{m}{i} \binom{n}{i}}{\binom{N}{i}} \quad (1)$$

where *m* and *n* are the number of proteins in the midbody and cilia proteomes, *k* is the number of overlapping proteins, and *N* is the size of the human proteome.

### Network Analysis

The functional network of human genes was prepared as described previously [Gray et al., 2009] and is searchable at: <http://www.functionalnet.org/humannet/>

As a measure of connectivity, we asked whether the proteins most strongly connected to midbody proteins in the gene network tended to be ciliary proteins using a method of Gaussian field label propagation [Mostafavi et al., 2008]. First, a bias score was assigned to each gene in the network: +1 to known midbody genes and  $k$  to all other genes. The value  $k$  is defined as:

$$k = \frac{n}{N} \quad (2)$$

where  $n$  is the number of midbody genes and  $N$  is the total number of genes in the network. The connectivity score was then computed as:

$$f = \arg \min_f \sum_i (f_i - y_i)^2 + \sum_i \sum_j w_{ij} (f_i - f_j)^2 \quad (3)$$

where  $W$  is an association matrix where entry  $w_{ij}$  is the network edge weight between genes  $i$  and  $j$ ,  $y_i$  is the bias node of  $i$ , and  $f$  is the vector of final scores where entry  $f_i$  indicates how strongly connected gene  $i$  is to the set of midbody genes. Each gene was rank-ordered by its final score as described previously [Lee et al., 2008b], and true positive and false positive rates were calculated as a function of rank. From the resultant ROC curve, the area under the ROC curve (AUC) was calculated. To obtain the P-value, the mean and standard deviation of AUCs obtained from 1000 random sets were calculated and a one-tailed Z-test was performed.

### C. elegans Strains and Assays

*C. elegans* strains were cultured in standard growth conditions on nematode growth medium with *Escherichia coli* strain OP50. Except where indicated, all worms were reared at 25°C. All assays were performed at 25°C. Temperature sensitive strains were maintained at 15°C with the exception of assays. Strains used were *lin-15B*(n765), *spd-1*(oj5), *zen-4*(or153), and *kyls104*(str-1p::gfp). *spd-1*(oj5) and *zen-4*(or153) strains expressing GFP behind the *str-1* promoter were made using standard mating techniques. All strains were provided by the *Caenorhabditis* Genetics Center.

Temperature sensitive *spd-1* and *zen-4* mutant strains were grown at 15°C. Embryos were shifted to the restrictive temperature of 25°C when they were between the twofold stage and L1. This timing allowed PRC1/MKLP-1 function to be disrupted shortly after the commencement of *daf-19* expression—seen to be from the twofold stage to early L2 [Swoboda et al., 2000]—while avoiding the early embryonic cell divisions. *spd-1* adult worms that had been raised at either the permissive or restrictive temperatures were assayed using chemotaxis plates set up in a variation on previous protocols [Troemel et al., 1997]. Briefly, adult worms were washed from OP50 fed plates using M9 buffer, and washed again with M9 until all bacteria were removed. Worms were then placed on the cen-

ter of the agar plate, equidistant from two circles—one soaked with 50  $\mu$ l of 100 mM ammonium acetate attractant, the other empty—and allowed to migrate for 1 hr, at which time the worms within each circle were counted. FITC filling was performed using standard protocols [Hedgecock et al., 1985], and the fluorescent intensity of confocal projections was measured using Image Pro Plus.

### Imaging of *C. elegans* Cilia and Neurons

Worms for cilia and neuron imaging were mounted on agar pads and anesthetized with sodium azide. Confocal images of the AWB neuron cilia and dye-filled cell bodies were obtained using a Zeiss LSM5 PASCAL microscope with 63x plan Apochromat (1.4na) or 100x Fluar (1.3na) oil-immersion objectives. Confocal optical slices were collected using Zeiss LSM5 software and three-dimensional reconstructions of the cilia were made using Imaris x64 4.5.2.

### Acknowledgment

The authors thank Phil Abitua for technical assistance. All strains were provided by the *Caenorhabditis* Genetics Center which is funded by the NIH/NCRR. This work was supported by grants to E.M.M. from the NSF, NIH, Welch Foundation (F-1515), Texas Institute for Drug and Diagnostic Development, and a Packard Fellowship; to R.H.G and S.G.B. from a VIDI/Aspasia grant from Dutch Organisation for Scientific Research (VIDI 91766354) and EU FP7 consortium SYSCILIA; and to J.B.W. from the NIH/NIGMS, The March of Dimes, The Burroughs Wellcome Fund, the American Asthma Foundation, and the Texas Advanced Research Program. J.B.W. is an Early Career Scientist of the Howard Hughes Medical Institute.

### References

- Albertson R, Cao J, Hsieh TS, Sullivan W. 2008. Vesicles and actin are targeted to the cleavage furrow via furrow microtubules and the central spindle. *J Cell Biol* 181: 777–790.
- Andersen JS, Wilkinson CJ, Mayor T, Mortensen P, Nigg EA, Mann M. 2003. Proteomic characterization of the human centrosome by protein correlation profiling. *Nature* 426: 570–574.
- Avidor-Reiss T, Maer AM, Koundakjian E, Polyanovsky A, Keil T, Subramaniam S, Zuker CS. 2004. Decoding cilia function: defining specialized genes required for compartmentalized cilia biogenesis. *Cell* 117: 527–539.
- Baldari CT, Rosenbaum J. 2010. Intraflagellar transport: it's not just for cilia anymore. *Curr Opin Cell Biol* 22: 75–80.
- Becker BE, Gard DL. 2006. Visualization of the cytoskeleton in *Xenopus* oocytes and eggs by confocal immunofluorescence microscopy. *Methods Mol Biol* 322: 69–86.
- Berman SA, Wilson NF, Haas NA, Lefebvre PA. 2003. A novel MAP kinase regulates flagellar length in *Chlamydomonas*. *Curr Biol* 13: 1145–1149.
- Burghoorn J, Dekkers MP, Rademakers S, de Jong T, Willemsen R, Jansen G. 2007. Mutation of the MAP kinase DYF-5 affects



- docking and undocking of kinesin-2 motors and reduces their speed in the cilia of *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 104: 7157–7162.
- Fariss RN, Molday RS, Fisher SK, Matsumoto B. 1997. Evidence from normal and degenerating photoreceptors that two outer segment integral membrane proteins have separate transport pathways. *J Comp Neurol* 387: 148–156.
- Finetti F, Paccani SR, Riparbelli MG, Giacomello E, Perinetti G, Pazour GJ, Rosenbaum JL, Baldari CT. 2009. Intraflagellar transport is required for polarized recycling of the TCR/CD3 complex to the immune synapse. *Nat Cell Biol* 11: 1332–1339.
- Frisch D, Farbman AI. 1968. Development of order during ciliogenesis. *Anat Rec* 162: 221–232.
- Gherman A, Davis EE, Katsanis N. 2006. The ciliary proteome database: an integrated community resource for the genetic and functional dissection of cilia. *Nat Genet* 38: 961–962.
- Glotzer M. 2005. The molecular requirements for cytokinesis. *Science* 307: 1735–1739.
- Glotzer M. 2009. The 3Ms of central spindle assembly: microtubules, motors and MAPs. *Nat Rev Mol Cell Biol* 10: 9–20.
- Gray RS, Abitua PB, Wlodarczyk BJ, Szabo-Rogers HL, Blanchard O, Lee I, Weiss GS, Liu KJ, Marcotte EM, Wallingford JB, et al. 2009. The planar cell polarity effector Fuz is essential for targeted membrane trafficking, ciliogenesis and mouse embryonic development. *Nat Cell Biol* 11: 1225–1232.
- Gromley A, Yeaman C, Rosa J, Redick S, Chen CT, Mirabelle S, Guha M, Sillibourne J, Doxsey SJ. 2005. Centriolin anchoring of exocyst and SNARE complexes at the midbody is required for secretory-vesicle-mediated abscission. *Cell* 123: 75–87.
- Hayes JM, Kim SK, Abitua PB, Park TJ, Herrington ER, Kitayama A, Grow MW, Ueno N, Wallingford JB. 2007. Identification of novel ciliogenesis factors using a new in vivo model for mucociliary epithelial development. *Dev Biol* 312: 115–130.
- Hedgecock EM, Culotti JG, Thomson JN, Perkins LA. 1985. Axonal guidance mutants of *Caenorhabditis elegans* identified by filling sensory neurons with fluorescein dyes. *Dev Biol* 111: 158–170.
- Horvath J, Fliegau M, Olbrich H, Kispert A, King SM, Mitchison H, Zariwala MA, Knowles MR, Sudbrak R, Fekete G, et al. 2005. Identification and analysis of axonemal dynein light chain 1 in primary ciliary dyskinesia patients. *Am J Respir Cell Mol Biol* 33: 41–47.
- Hubbard TJ, Aken BL, Ayling S, Ballester B, Beal K, Bragin E, Brent S, Chen Y, Clapham P, Clarke L, et al. 2009. Ensembl 2009. *Nucleic Acids Res* 37: D690–D697.
- Inglis PN, Boroevich KA, Leroux MR. 2006. Piecing together a cilium. *Trends Genet* 22: 491–500.
- Jiang W, Jimenez G, Wells NJ, Hope TJ, Wahl GM, Hunter T, Fukunaga R. 1998. PRC1: a human mitotic spindle-associated CDK substrate protein required for cytokinesis. *Mol Cell* 2: 877–885.
- Keller LC, Romijn EP, Zamora I, Yates JR, III, Marshall WF. 2005. Proteomic analysis of isolated chlamydomonas centrioles reveals orthologs of ciliary-disease genes. *Curr Biol* 15: 1090–1098.
- Kieserman EK, Glotzer M, Wallingford JB. 2008. Developmental regulation of central spindle assembly and cytokinesis during vertebrate embryogenesis. *Curr Biol* 18: 116–123.
- Kim J, Lee JE, Heynen-Genel S, Suyama E, Ono K, Lee K, Ideker T, Aza-Blanc P, Gleeson JG. 2010. Functional genomic screen for modulators of ciliogenesis and cilium length. *Nature* 464: 1048–1051.
- Kim JC, Ou YY, Badano JL, Esmail MA, Leitch CC, Friedrich E, Beales PL, Archibald JM, Katsanis N, Rattner JB, et al. 2005. MKKS/BBS6, a divergent chaperonin-like protein linked to the obesity disorder Bardet-Biedl syndrome, is a novel centrosomal component required for cytokinesis. *J Cell Sci* 118: 1007–1020.
- Kurasawa Y, Earnshaw WC, Mochizuki Y, Dohmae N, Todokoro K. 2004. Essential roles of KIF4 and its binding partner PRC1 in organized central spindle midzone formation. *EMBO J* 23: 3237–3248.
- Lee CJ, Kieserman E, Gray RS, Park TJ, Wallingford JB. 2008a. Whole-mount fluorescence immunocytochemistry on *Xenopus* embryos. *Cold Spring Harb Protoc*.
- Lee I, Lehner B, Crombie C, Wong W, Fraser AG, Marcotte EM. 2008b. A single gene network accurately predicts phenotypic effects of gene perturbation in *Caenorhabditis elegans*. *Nat Genet* 40: 181–188.
- Li JB, Gerdes JM, Haycraft CJ, Fan Y, Teslovich TM, May-Simera H, Li H, Blacque OE, Li L, Leitch CC, et al. 2004. Comparative genomics identifies a flagellar and basal body proteome that includes the BBS5 human disease gene. *Cell* 117: 541–552.
- Li Z, Lee I, Moradi E, Hung NJ, Johnson AW, Marcotte EM. 2009. Rational extension of the ribosome biogenesis pathway using network-guided genetics. *PLoS Biol* 7: e1000213.
- Liu Q, Tan G, Levenkova N, Li T, Pugh EN, Jr., Rux JJ, Speicher DW, Pierce EA. 2007. The proteome of the mouse photoreceptor sensory cilium complex. *Mol Cell Proteomics* 6: 1299–1317.
- Mahjoub MR, Qasim Rasi M, Quarmby LM. 2004. A NIMA-related kinase, Fa2p, localizes to a novel site in the proximal cilia of *Chlamydomonas* and mouse kidney cells. *Mol Biol Cell* 15: 5172–5186.
- Marshall WF. 2008. The cell biological basis of ciliary disease. *J Cell Biol* 180: 17–21.
- Mitchell B, Jacobs R, Li J, Chien S, Kintner C. 2007. A positive feedback mechanism governs the polarity and motion of motile cilia. *Nature* 447: 97–101.
- Mostafavi S, Ray D, Warde-Farley D, Grouios C, Morris Q. 2008. GeneMANIA: a real-time multiple association network integration algorithm for predicting gene function. *Genome Biol* 9 (Suppl 1): S4.
- Mukhopadhyay S, Lu Y, Qin H, Lanjuin A, Shaham S, Sengupta P. 2007. Distinct IFT mechanisms contribute to the generation of ciliary structural diversity in *C. elegans*. *EMBO J* 26: 2966–2980.
- Mukhopadhyay S, Lu Y, Shaham S, Sengupta P. 2008. Sensory signaling-dependent remodeling of olfactory cilia architecture in *C. elegans*. *Dev Cell* 14: 762–774.
- Murayama T, Toh Y, Ohshima Y, Koga M. 2005. The dyf-3 gene encodes a novel protein required for sensory cilium formation in *Caenorhabditis elegans*. *J Mol Biol* 346: 677–687.
- Nachury MV, Loktev AV, Zhang Q, Westlake CJ, Peranen J, Merdes A, Slusarski DC, Scheller RH, Bazan JF, Sheffield VC, et al. 2007. A core complex of BBS proteins cooperates with the GTPase Rab8 to promote ciliary membrane biogenesis. *Cell* 129: 1201–1213.
- Neef R, Gruneberg U, Kopajtich R, Li X, Nigg EA, Sillje H, Barr FA. 2007. Choice of Plk1 docking partners during mitosis and cytokinesis is controlled by the activation state of Cdk1. *Nat Cell Biol* 9: 436–444.
- Neugebauer JM, Amack JD, Peterson AG, Bisgrove BW, Yost HJ. 2009. FGF signalling during embryo development regulates cilia length in diverse epithelia. *Nature* 458: 651–654.
- O'Connell KF, Leys CM, White JG. 1998. A genetic screen for temperature-sensitive cell-division mutants of *Caenorhabditis elegans*. *Genetics* 149: 1303–1321.

- Pan J, You Y, Huang T, Brody SL. 2007. RhoA-mediated apical actin enrichment is required for ciliogenesis and promoted by Foxj1. *J Cell Sci* 120: 1868–1876.
- Park TJ, Haigo SL, Wallingford JB. 2006. Ciliogenesis defects in embryos lacking inturned or fuzzy function are associated with failure of planar cell polarity and Hedgehog signaling. *Nat Genet* 38: 303–311.
- Park TJ, Mitchell BJ, Abitua PB, Kintner C, Wallingford JB. 2008. Dishevelled controls apical docking and planar polarization of basal bodies in ciliated epithelial cells. *Nat Genet* 40: 871–879.
- Perkins LA, Hedgecock EM, Thomson JN, Culotti JG. 1986. Mutant sensory cilia in the nematode *Caenorhabditis elegans*. *Dev Biol* 117: 456–487.
- Potapova TA, Daum JR, Pittman BD, Hudson JR, Jones TN, Sattinover DL, Stukenberg PT, Gorbsky GJ. 2006. The reversibility of mitotic exit in vertebrate cells. *Nature* 440: 954–958.
- Qin H, Wang Z, Diener D, Rosenbaum J. 2007. Intraflagellar transport protein 27 is a small G protein involved in cell-cycle control. *Curr Biol* 17: 193–202.
- Ren J, Liu Z, Gao X, Jin C, Ye M, Zou H, Wen L, Zhang Z, Xue Y, Yao X. 2009. MiCroKit 3.0: an integrated database of midbody, centrosome and kinetochore. *Nucleic Acids Res* 38: D155–160.
- Shah AS, Farnen SL, Moninger TO, Businga TR, Andrews MP, Bugge K, Searby CC, Nishimura D, Brogden KA, Kline JN, et al. 2008. Loss of Bardet-Biedl syndrome proteins alters the morphology and function of motile cilia in airway epithelia. *Proc Natl Acad Sci USA* 105: 3380–3385.
- Sharma N, Berbari NF, Yoder BK. 2008. Ciliary dysfunction in developmental abnormalities and diseases. *Curr Top Dev Biol* 85: 371–427.
- Sive HL, Grainger RM, Harland RM. 2000. *Early Development of Xenopus laevis: A Laboratory Manual*. Cold Spring Harbor, N.Y.: Cold Spring Harbor Press.
- Skop AR, Liu H, Yates J, III, Meyer BJ, Heald R. 2004. Dissection of the mammalian midbody proteome reveals conserved cytokinesis mechanisms. *Science* 305: 61–66.
- Sorokin SP. 1968. Reconstructions of centriole formation and ciliogenesis in mammalian lungs. *J Cell Sci* 3: 207–230.
- Spektor A, Tsang WY, Khoo D, Dynlacht BD. 2007. Cep97 and CP110 suppress a cilia assembly program. *Cell* 130: 678–690.
- Steinman RM. 1968. An electron microscopic study of ciliogenesis in developing epidermis and trachea in the embryo of *Xenopus laevis*. *Am J Anat* 122: 19–55.
- Swoboda P, Adler HT, Thomas JH. 2000. The RFX-type transcription factor DAF-19 regulates sensory neuron cilium formation in *C. elegans*. *Mol Cell* 5: 411–421.
- Troemel ER, Kimmel BE, Bargmann CI. 1997. Reprogramming chemotaxis responses: sensory neurons define olfactory preferences in *C. elegans*. *Cell* 91: 161–169.
- Verbrugghe KJ, White JG. 2004. SPD-1 is required for the formation of the spindle midzone but is not essential for the completion of cytokinesis in *C. elegans* embryos. *Curr Biol* 14: 1755–1760.
- Yang J, Li T. 2005. The ciliary rootlet interacts with kinesin light chains and may provide a scaffold for kinesin-1 vesicular cargos. *Exp Cell Res* 309: 379–389.
- Zuo X, Guo W, Lipschutz JH. 2009. The exocyst protein Sec10 is necessary for primary ciliogenesis and cystogenesis in vitro. *Mol Biol Cell* 20: 2522–2529.