

Bacteriophages use an expanded genetic code on evolutionary paths to higher fitness

Michael J Hammerling, Jared W Ellefson, Daniel R Boutz, Edward M Marcotte, Andrew D Ellington & Jeffrey E Barrick*

Bioengineering advances have made it possible to fundamentally alter the genetic codes of organisms. However, the evolutionary consequences of expanding an organism's genetic code with a noncanonical amino acid are poorly understood. Here we show that bacteriophages evolved on a host that incorporates 3-iodotyrosine at the amber stop codon acquire neutral and beneficial mutations to this new amino acid in their proteins, demonstrating that an expanded genetic code increases evolvability.

The evolution of life on Earth has been dictated by the underlying constancy of a nearly universal genetic code with twenty amino acids^{1,2}. However, examples of natural genetic codes that have been functionally expanded with a twenty-first amino acid³ and the multitude of known post-translational protein modifications suggest that, though aspects of the genetic code have been optimized by evolution^{4,5}, it does not provide the necessary chemical diversity to best perform all functions of potential benefit to organisms. Technologies exist to augment genetic codes with a noncanonical amino acid (ncAA) by introducing an orthogonal aminoacyl-tRNA synthetase (aaRS) and a cognate tRNA recognizing the amber stop codon^{6,7}. We hypothesized that organisms given the ability to encode a twenty-first amino acid would evolve to use this new chemical building block on mutational pathways to higher fitness.

Several challenges can arise when attempting to evolve an organism with a newly expanded genetic code, and the directed evolution of even single proteins with ncAAs has been limited to date⁸. The organism of interest must first survive the globally disruptive change in how its genomic information is decoded into its proteome². Next, the reassigned codon must be translated with sufficient efficiency and fidelity for substitutions of the new codon to be beneficial, which is less likely if translating this codon sometimes results in truncated proteins⁹ or ambiguous amino acid incorporation¹⁰. Evolution may also need to proceed for many generations to observe ncAA substitutions because only a small fraction of mutations will result in changes to the reassigned codon. Finally, one must circumvent rejection of the reengineered genetic code in cases where mutations that lead to the ncAA no longer being incorporated into proteins are highly advantageous to the host organism¹¹.

Bacteriophage T7 has been used as a model organism for studies of molecular genetics^{12,13}, experimental evolution^{14,15} and synthetic biology¹⁶. We evolved this bacterial virus in a constant translational environment with an expanded genetic code by passaging it on an RF0 IodoY *Escherichia coli* host that efficiently incorporates 3-iodotyrosine at amber stop codons owing to the deletion of protein release factor 1 (RF1) and addition of an engineered aaRS-tRNA pair^{17,18} (**Supplementary Results, Supplementary Figs. 1–3**). Populations of wild-type T7 (WT)¹² and a hypermutator variant

($\Delta 2$)¹⁹ were propagated on cultures of actively growing *E. coli* hosts by incubating them together until there was visible bacterial lysis due to viral replication and then transferring a fraction of this mixture to a new *E. coli* culture (**Fig. 1a**). Lysis time decreased for both WT (from ~100 min to ~35 min) and $\Delta 2$ (from ~120 min to ~45 min) populations over 50 serial transfers, indicating that they evolved higher fitness on the RF0 IodoY *E. coli* host.

An organism with a newly expanded genetic code is subject to new constraints on codon usage and may have new opportunities for adaptive mutations. The evolutionary response could include: (i) compensatory mutations that restore protein termination in cases where readthrough of a reassigned stop codon is deleterious, (ii) genetic drift to reassigned codons at positions where ncAA substitutions are selectively neutral or (iii) mutations to reassigned codons that are potentially beneficial in ways that would not be possible with a canonical amino acid. Deleterious mutations are highly unlikely to contribute to evolution in our experiment owing to the extremely large effective population size ($\sim 10^7$ – 10^8 phages transferred each passage).

If amber codons evolve in important or essential genes, these mutations are expected to result in addiction, such that an organism requires an alternative genetic code for viability. To test for this level of dependence, we titered the evolved T7 populations on three *E. coli* hosts, which either terminate translation (BL21(DE3)) or incorporate 3-iodotyrosine (RF0 IodoY) or tyrosine (RF0 Tyr) at the amber codon (**Fig. 1b**). All of the populations produced statistically indistinguishable numbers of phage plaques on RF0 IodoY and RF0 Tyr cells, but one WT and three $\Delta 2$ populations produced significantly ($P \leq 0.01$) fewer plaques on the BL21(DE3) host (**Fig. 1c**), indicating that some phages in these populations had evolved amber codons inside key genes.

To determine exactly what genetic changes had taken place, we sequenced DNA samples isolated from four evolved T7 populations. Despite an approximately tenfold difference in mutation rates¹⁹, the base substitution spectra in $\Delta 2$ and WT populations were similar (**Supplementary Fig. 4**). However, the overall character of genome evolution shifted from directional selection in the wild-type populations (dN/dS = 1.58) to purifying selection in the hypermutator populations (dN/dS = 0.57). Deletions overlapping gene 0.7 that are known to be beneficial^{13,15} were present, in aggregate, in ~100% of every sequenced population (**Supplementary Fig. 5 and Supplementary Table 1**), and there was a deletion overlapping two hypothetical genes in one population (**Supplementary Fig. 6**).

Among the point mutations present in these populations (**Fig. 1d**), we observed two substitutions of amber codons that reached high frequencies: one in T7 RNA polymerase (RNAP) and the other in the T7 type II holin. Patching random phage isolates from the




Figure 1 | Genome evolution of a bacterial virus with a newly expanded genetic code. (a) In the evolution experiment, six populations each of WT and $\Delta 2$ bacteriophage T7 were passaged on a release factor 1 knockout (RFO) *E. coli* host strain carrying an orthogonal tRNA and aaRS system that recodes the amber stop codon to the non-natural amino acid 3-iodotyrosine. **(b)** *E. coli* host strains used to test whether evolved phages required amber suppression by 3-iodotyrosine or tyrosine to replicate. **(c)** Relative titer (number of plaques) formed by each evolved bacteriophage population on the host strains in **b**. Values are the average of three technical replicates with error bars that represent 95% confidence limits. *P* values for reduced phage number on the alternative hosts compared to RFO IodoY are for Student's *t*-tests assuming equal variance and using the Bonferroni correction for multiple testing. **(d)** Locations of point mutations with frequencies $\geq 5\%$ observed in metagenomic DNA sequencing data for the four bacteriophage populations boxed in **c**. Red open reading frames encode T7 proteins with known functions¹³. Mutations of interest in regions highlighted in yellow are discussed in the text and following figures. There were roughly 50 and 400 mutations present with frequencies $\geq 1\%$ in the evolved WT and $\Delta 2$ T7 populations, respectively.

respective populations confirmed that each new amber codon was associated with a defect in forming plaques on the BL21(DE3) normal genetic code *E. coli* host. As anticipated, we also observed compensation for readthrough of phage proteins ending in the amber codon. Notably, mutations appeared to restore T7 exonuclease (gene 6) termination in close to 100% of all four sequenced populations (Supplementary Fig. 7).

T7 RNAP is an essential protein that is responsible for transcription of bacteriophage genes¹³. The Trp88-to-amber mutation in RNAP (Supplementary Fig. 8a) probably has little effect on phage fitness on the RFO IodoY host. Of 37 base substitutions observed in this gene with frequencies $\geq 1\%$ across all populations, 21 were synonymous. Thus, the overall molecular signature of selection within RNAP is highly purifying ($dN/dS = 0.22$). So, even nonsynonymous changes such as this amber mutation most likely were tolerated only because they did not alter protein function. Furthermore, only one of these 37 mutations was present in a WT population, where fewer neutral mutations are expected. Finally, Trp88 is located at a surface-exposed site in the structure of RNAP²⁰ where 3-iodotyrosine seems to be accommodated without altering the surrounding molecular structure (Supplementary Fig. 8b).

In contrast, several lines of evidence indicate that the Tyr39-to-amber substitution in the T7 type II holin (gene 17.5) is beneficial to phage fitness (Fig. 2a). Holins regulate lysis timing by forming pores in the bacterial membrane through which lysozyme degrades the cell wall²¹, and optimizing lysis time is critical to maximizing phage progeny^{22,23}. We found a higher density of point mutations with frequencies $\geq 1\%$ within gene 17.5 than in the genome at large (binomial test; $P = 1.5 \times 10^{-9}$). Furthermore, all 22 of these base substitutions in the type II holin were nonsynonymous (Supplementary Table 2), and nearly 100% of phages in each WT and $>50\%$ in each $\Delta 2$ population have a mutation somewhere in gene 17.5. Together these

observations indicate that type II holin activity was under strong directional selection during the evolution experiment.

The amber mutation in the T7 type II holin occurs with the highest frequency (53%) of any coding base substitution in this population and with the highest frequency of any gene 17.5 mutation in any sequenced phage population. This result is even more notable given that it is in a wild-type rather than a hypermutator background. As expected for a highly beneficial mutation, phages with the amber lyse RFO IodoY cells more completely at earlier times than phages from the same population without it, and this difference is not found on the RFO Tyr host, where tyrosine incorporation at the amber codon restores the normal type II holin sequence (Supplementary Fig. 9).

To directly test whether the 3-iodotyrosine substitution in the type II holin protein was beneficial to phage fitness, we isolated spontaneous rescue mutants of a phage with this amber that could plaque on normal genetic code cells. Only tryptophan and tyrosine mutants were obtained, suggesting that position 39 of the holin is constrained to aromatic amino acids (Fig. 2b). The evolved isolate coding 3-iodotyrosine was more fit than the rescue mutants with tyrosine or tryptophan at this position when they were competed in co-culture under the conditions of the evolution experiment (Fig. 2c). This outcome held for a wide range of transfer volumes that altered the initial multiplicity of infection at each cycle (Supplementary Fig. 10).

We have shown that access to an expanded genetic code increased the evolvability of an organism by characterizing a new beneficial mutation that was only possible with 3-iodotyrosine incorporation into proteins. The readiness with which bacteriophage T7 adapted to and used a newly expanded genetic code with a ncAA was unexpected. This result establishes that current technologies can effectively thaw Crick's 'frozen accident', opening vast new sequence spaces and new




Figure 2 | Beneficial amber mutation in the T7 holin II protein.

(a) The frequency of an amber mutation resulting in a substitution from tyrosine to 3-iodotyrosine in the T7 type II holin (gene 17.5) can explain the reduced titer of population WT L6 on the standard genetic code host (Fig. 1c). Signatures of molecular evolution described in the text strongly suggest that this mutation increases phage fitness on the RF0 lodoY *E. coli* host. (b) Rescue mutants of an evolved WT L6 phage isolate with this amber mutation were obtained by plating on *E. coli* BL21(DE3) hosts with the standard genetic code and isolating fast-growing plaques. Of the eight possible neighboring codons accessible by single-base mutations that code for amino acids, only tyrosine and tryptophan mutants were observed. The numbers of rescue mutants found with each codon are indicated at the end of the arrows. (c) The evolved phage isolate with the amber codon (TAG) at position 39 of the holin protein was competed against rescue mutants with a tyrosine (TAC) or tryptophan (TGG) at that position. In each competition, lysates of the two phage preconditioned separately were mixed and then passaged on RF0 lodoY cells for three transfers. Sanger sequencing traces at this position show that the amber codon increases in frequency relative to the alternative, indicating that 3-iodotyrosine confers a fitness benefit compared to both the ancestral amino acid and a chemically similar amino acid tolerated at this position.

chemistries for investigation. These capabilities will continue to improve as the genomes of organisms are engineered to have one or more blank codons that can be reassigned to ncAAs without disrupting existing proteins²⁴. Longer evolution experiments are likely to result in codon substitutions that addict organisms to the unique functionality of an introduced ncAA for survival. These may be useful for semantic biocontainment of genetically modified organisms²⁵ and for whole-genome evolutionary approaches for further improving ncAA incorporation. Finally, this general research strategy makes it possible to experimentally address questions about the optimality of the natural genetic code and to discover unexpected utility for genetically encoded ncAAs by the directed evolution of proteins and entire organisms.

Received 11 August 2013; accepted 19 December 2013; published online 2 February 2014

Methods

Methods and any associated references are available in the [online version of the paper](#).

Accession codes. National Center for Biotechnology Information Short Read Archive: Genome resequencing data for bacteriophage

clones and populations have been deposited under accession number SRP021199.

References

- Crick, F.H.C. *J. Mol. Biol.* **38**, 367–379 (1968).
- Knight, R.D., Freeland, S.J. & Landweber, L.F. *Nat. Rev. Genet.* **2**, 49–58 (2001).
- Ambrogelly, A., Palioura, S. & Söll, D. *Nat. Chem. Biol.* **3**, 29–35 (2007).
- Freeland, S.J. & Hurst, L.D. *J. Mol. Evol.* **47**, 238–248 (1998).
- Itzkovitz, S. & Alon, U. *Genome Res.* **17**, 405–412 (2007).
- Liu, C.C. & Schultz, P.G. *Annu. Rev. Biochem.* **79**, 413–444 (2010).
- Davis, L. & Chin, J.W. *Nat. Rev. Mol. Cell Biol.* **13**, 168–182 (2012).
- Brustad, E.M. & Arnold, F.H. *Curr. Opin. Chem. Biol.* **15**, 201–210 (2011).
- Wang, K., Neumann, H., Peak-Chew, S.Y. & Chin, J.W. *Nat. Biotechnol.* **25**, 770–777 (2007).
- Bacher, J.M., Hughes, R.A., Tze-Fei Wong, J. & Ellington, A.D. *Trends Ecol. Evol.* **19**, 69–75 (2004).
- Ohtake, K. *et al. J. Bacteriol.* **194**, 2606–2613 (2012).
- Dunn, J.J. & Studier, F.W. *J. Mol. Biol.* **166**, 477–535 (1983).
- Molineux, I.J. *The Bacteriophages 277–301* (Oxford UP, 2006).
- Bull, J.J. *et al. Genetics* **147**, 1497–1507 (1997).
- Bull, J.J., Badgett, M.R., Rokyta, D. & Molineux, I.J. *J. Mol. Evol.* **57**, 241–248 (2003).
- Chan, L.Y., Kosuri, S. & Endy, D. *Mol. Syst. Biol.* **1**, 2005.0018 (2005).
- Sakamoto, K. *et al. Structure* **17**, 335–344 (2009).
- Mukai, T. *et al. Biochem. Biophys. Res. Commun.* **411**, 757–761 (2011).
- Tabor, S. & Richardson, C.C. *J. Biol. Chem.* **264**, 6447–6458 (1989).
- Cheetham, G.M. & Steitz, T.A. *Science* **286**, 2305–2309 (1999).
- Wang, I.N., Smith, D.L. & Young, R. *Annu. Rev. Microbiol.* **54**, 799–825 (2000).
- Wang, N., Dykhuizen, D. & Slobodkin, L. *Evol. Ecol.* **10**, 545–558 (1996).
- Heineman, R.H., Bull, J.J. & Molineux, I.J. *Mol. Biol. Evol.* **26**, 1289–1298 (2009).
- Lajoie, M.J. *et al. Science* **342**, 357–360 (2013).
- Moe-Behrens, G.H.G., Davis, R. & Haynes, K.A. *Front. Microbiol.* **4**, 5 (2013).

Acknowledgments

We thank J. Bull, I. Molineux, R. Hughes, C. Barnhart, D. Deatherage, R. Alnahhas, M. Schmerer, A. Miklos, A. Meyer and A. Maranhão (all at the University of Texas at Austin) for plasmids, strains, advice and technical assistance. The RF0 lodoY strain was provided by RIKEN and by the Targeted Proteins Research Program, the Ministry of Education, Culture, Sports, Science and Technology, Japan. We acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing HPC resources. This research was supported by the US National Institutes of Health (NIH) (R00-GM087550 to J.E.B.), the US National Science Foundation (NSF) BEACON Center for the Study of Evolution in Action (DBI-0939454 to J.E.B.), the US Army Research Office (W911NF-12-1-0390 to J.E.B. and E.M.M.), the US National Security Science and Engineering Faculty (FA9550-10-1-01-69 to A.D.E.), the US Defense Advanced Research Project Agency (HR-0011-12-C-0066 to A.D.E.) and the NSF (MCB-0943383 to A.D.E.). E.M.M. also acknowledges funding from the NIH, the NSF, the Cancer Prevention Research Institute of Texas and the Welch Foundation (F1515).

Author contributions

M.J.H., J.W.E., J.E.B. and A.D.E. conceived the study. M.J.H. performed evolution experiments. J.E.B. and M.J.H. analyzed sequencing data. M.J.H. and J.W.E. created RF0 Tyr and characterized phage lysis times. D.R.B. and E.M.M. analyzed proteomics data. J.E.B. performed statistical analyses. J.E.B., M.J.H., J.W.E. and D.R.B. created figures and wrote the manuscript. All of the authors designed experiments and edited the manuscript.

Competing financial interests

The authors declare no competing financial interests.

Additional information

Supplementary information is available in the [online version of the paper](#). Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>. Correspondence and requests for materials should be addressed to J.E.B.

ONLINE METHODS

***E. coli* host strains.** RF0 IodoY¹⁸ was constructed from BL21(DE3) *E. coli* (Invitrogen) by creating a chloramphenicol-resistant F plasmid containing several key *E. coli* genes with recoded stop codons, knocking out RF1 in the chromosome and adding a second plasmid containing a modified *Methanococcus jannaschii* tyrosyl-tRNA synthetase and UAG-decoding tRNA for specifically incorporating 3-iodotyrosine at the amber stop codon¹⁷. Cultures of RF0 IodoY were supplemented with 20 µg/mL chloramphenicol and 10 µg/mL gentamycin. RF0 Tyr was generated by adding a plasmid created by replacing the gentamycin resistance marker in the RF0 IodoY aaRS-tRNA plasmid with a spectinomycin resistance gene and by replacing the 3-iodotyrosine aaRS with the wild-type *M. jannaschii* tyrosyl-tRNA synthetase (GenBank accession code NP_247363.1). RF0 Tyr cultures were supplemented with 60 µg/mL spectinomycin and 10 µg/mL gentamycin. All of these strains contain a copy of T7 RNAP in the *E. coli* host genome under control of the *lac* UV5 promoter. This promoter was not induced during any of our experiments, and background expression of this host copy of T7 RNAP was apparently low enough that phages with an amber stop codon in their copy of the T7 RNAP gene could not plaque on BL21(DE3) without amber suppression.

Phage evolution. Wild-type T7 and hypermutator Δ2 phage ancestors were obtained from I.J. Molineux (University of Texas at Austin). A single plaque of each was isolated on BL21(DE3) cells. Frozen aliquots of RF0 IodoY cells were thawed; inoculated into 50-mL Erlenmeyer flasks containing 10 mL Lysogeny Broth (LB), appropriate antibiotics and 250 µM 3-iodotyrosine (Sigma-Aldrich); and grown to mid-exponential phase (OD₆₀₀ ~0.5) at 37 °C with orbital shaking at 120 r.p.m. Six lines each of wild-type and mutator phages were initiated by seeding separate *E. coli* cultures with 1 µL of clonal phage lysate. Phages were allowed to replicate until all of the flasks showed visible lysis by clearing of their turbidity, a period of time that varied from ~100–120 min at the beginning of the evolution experiment to ~35–45 min at the end. After lysis, 1 µL of each independent culture was transferred to fresh RF0 IodoY cells that had been revived from the same freezer stock and grown to mid-exponential phase. This transfer volume yields a low initial multiplicity of infection (MOI) of ~0.05–0.1. Evolved phage populations were periodically archived at –80 °C in LB with 15% (v/v) glycerol added.

Genome sequencing and analysis. Phage DNA from each line was isolated using a CsCl₂ gradient and sequenced using standard methods on an Illumina MiSeq instrument at the Genome Sequencing and Analysis Facility (University of Texas at Austin). Adaptor sequences were trimmed from the resulting 250-base paired-end reads using FLEXBAR²⁶ (version 2.32).

We created updated reference sequences for the ancestral T7 wild-type and Δ2 hypermutator phages by comparing the reads from each sample to the published T7 genome sequence¹² (GenBank NC_001604.1) using breseq (version 0.23; <http://barricklab.org/breseq/>) in consensus mode. The wild-type ancestor had only a single difference from the published T7 sequence: an insertion of a single A base after position 1895 within gene 0.6B. The Δ2 ancestor had 112 genetic differences relative to the wild-type ancestor. These differences included 107 base substitutions, two single-nucleotide insertions, two single-nucleotide deletions and the in-frame 6-bp deletion in the exonuclease domain of T7 DNA polymerase that results in the hypermutator phenotype¹⁹. Complete lists of the genetic differences between the Δ2 and WT ancestors and the published T7 genome are available (**Supplementary Data Set 1**).

Mutations were predicted from the metagenomic sequencing data for the evolved populations using breseq in two stages. The first stage used breseq with both polymorphism and junction prediction enabled to predict large-scale structural variants. From these results, we estimated the fraction of evolved populations containing large deletions by manually examining regions with anomalous read coverage and by counting reads that mapped across new and ancestral junction sequences. The second analysis stage used breseq in polymorphism prediction mode with no prediction of new sequence junctions. This output was used to predict point mutation variants with frequencies ≥1% in each population (**Supplementary Data Set 1**). Calculated dN/dS ratios accounted for codon abundance and assumed an equal probability of all base substitutions.

Evolved phage population titers. Frozen stocks were first thawed, diluted and spotted on RF0 IodoY to determine the approximate titer of each stock. Then, 100 µL of an appropriate dilution was added to 3 mL of melted soft agar

supplemented with appropriate antibiotics and also 250 µM 3-iodotyrosine for RF0 IodoY. Three hundred microliters of *E. coli* culture at an optical density of ~1.0 was immediately added, and soft agar was poured across the surface of LB agar plates. Plates were incubated at 37 °C for 5 h, at which time the number of plaques on each host was counted.

Patching assays. Phage dilutions were plated on the permissive RF0 IodoY host at 37 °C and stored at 4 °C overnight after clearly discernable plaques were obtained. The next day, *E. coli* host strains to be tested were inoculated into soft agar containing appropriate supplements and poured over LB plates. Isolated plaques were then picked with a pipette tip and resuspended in 1 mL of LB, and 1 µL of this dilution was spotted on each host strain. Plates were incubated at 37 °C, and growth was assessed after 5 h. Lysates of a collection of plaques with and without the amber suppressor-dependent phenotype were created, and the allelic state of the amber-containing gene in each phage isolate was determined via PCR and Sanger sequencing.

Molecular modeling. PyMOL (Version 1.5.0.4; Schrödinger, LLC) with the SwissSideChain plugin²⁷ was used to model and visualize the 3-iodotyrosine substitution in the structure of RNA polymerase (Protein Data Bank code 1QLN)²⁰.

Lysis assays. RF0 IodoY and RF0 Tyr cells were revived and grown with 250 µM 3-iodotyrosine and appropriate antibiotics. When cultures reached an OD₆₀₀ of ~0.5, 200-µL aliquots were transferred into a clear-bottom 96-well plate and lysates of individual phage stocks, which had been previously titered on RF0 IodoY cells, were added to each well at a dilution that gave an MOI of ~5. These phages had known allelic states of the gene 17.5 amber mutation, verified by patching and sequencing. The microplate was incubated at 37 °C with OD₆₀₀ readings taken every minute, preceded by 10 s of shaking at intensity 4, in a PowerWave 340 microplate spectrophotometer (BioTek). Lysis curves were corrected for the time it took to aliquot each separate phage and to begin making measurements in the plate reader (<10 min).

Amber rescue mutants. Lysate was created from an evolved WT L6 phage clone that was isolated and confirmed to have the amber mutation in the type II holin gene by patching, PCR and Sanger sequencing, as above. For most rescue mutants, multiple independent cultures of RF0 IodoY were inoculated with ~10⁴ phage from this stock and again allowed to lyse completely, and 10³–10⁵ dilutions were plated on BL21(DE3) cells in soft agar as above. Other rescue mutants were isolated by plating from this stock directly on BL21(DE3) cells. In each case, mutations of the amber to nontermination codons were identified as the fastest-forming plaques, inoculated into RF0 IodoY cells to obtain lysate and genotyped via PCR and Sanger sequencing.

Competitions. Phages to be competed were revived from frozen stocks through two serial transfers in RF0 IodoY cells under the same conditions as the evolution experiment. After complete lysis of the second cultures, equal volumes of each phage were mixed and inoculated into new RF0 IodoY cultures to initiate competitions at 0.01×, 0.1×, 1× or 10× the transfer dilution volume used in the evolution experiment to test the effect of different initial phage MOI. Competitions proceeded for three serial transfers at their respective dilutions, and a sample of phage lysate was saved at the end of each infection cycle. After passaging was complete, whole-phage PCR was performed with 1 µL of the mixed lysate as template for the region of interest. Changes in the heights of peaks corresponding to each allele in Sanger sequencing traces for these PCR products over multiple passages were used to infer the more-fit phage genotype.

GFP emission assay. RF0 IodoY cells were transformed with pET21 plasmids containing sfGFP variants under the control of the *lac* operator and a strong T7 promoter. pET21 sfGFP (WT) contained a version of superfolder GFP (sfGFP) as a positive control. pET21 sfGFP (Y66→TAG) contained an sfGFP variant with an amber codon mutation replacing Y66 in the fluorophore. Individual transformants were isolated and grown overnight with appropriate antibiotics and nCAA supplements. The next day, saturated cultures were diluted 100-fold into the same medium and allowed to reach mid-log phase, at which time cultures were induced with 500 µM IPTG. Samples were taken 150 min later and analyzed on a Tecan Safire plate reader. An emission scan with excitation at 480 nm was performed to identify the peak emission wavelength for each

sample. RF0 IodoY pET21 GFP(Y66→TAG) was found to emit with a peak at 524 nm, which is 8 nm longer than the WT sfGFP peak of 516 nm and in accordance with previous studies of GFP with 3-iodotyrosine incorporated at this position²⁸.

Proteomics. RF0 IodoY *E. coli* cells cultured in LB supplemented with 3-iodotyrosine and appropriate antibiotics were pelleted by centrifugation and resuspended in 50 mM Tris (pH 8.0) followed by the addition of 2,2,2-trifluoroethanol (TFE) to a concentration of 50% (v/v) to solubilize the cell membrane and denature proteins. Samples were then reduced with 5 mM dithiothreitol (DTT) for 45 min at 56 °C and alkylated with 15 mM iodoacetamide for 30 min. Excess iodoacetamide was quenched by the addition of another 15 mM DTT. Samples were diluted tenfold in 50 mM Tris (pH 8.0) with 4 mM CaCl₂ to a final concentration of 5% TFE (v/v) before proteolytic digestion with trypsin for 4 h at 37 °C.

Alternatively, RF0 IodoY *E. coli* cells were lysed by a T7 phage isolate from evolved population WT L6 with the amber substitution in the type II holin (gene 17.5), or BL21(DE3) *E. coli* cells were lysed by wild-type T7. The insoluble pellet from each lysed sample was resuspended in SDS-PAGE loading buffer, and proteins were separated on a 4–20% acrylamide gradient gel. Gel slices corresponding to 25–17 kDa, 17–12 kDa and <12 kDa were cut and subjected to in-gel digestion (8 h at 37 °C) as previously described²⁹ but with endoproteinase GluC (New England BioLabs) instead of trypsin. Following digestion, peptides were eluted from the gel, dried by speedvac, resuspended in 50 mM Tris (pH 8.0) with 4 mM CaCl₂ and digested with trypsin as described above.

Peptides resulting from both in-gel and in-solution digestions were desalted on Hypersep C18 spin columns (Thermo Scientific) and analyzed by nano LC-MS/MS on a Dionex Ultimate 3000 nanoRSLC system coupled to an LTQ-Orbitrap Velos Pro mass spectrometer (Thermo Scientific). Data-dependent ion selection was activated with parent ion scans (MS1) collected at 100,000 (FWHM) resolution. Ions with charge > +1 were selected for collision-induced

dissociation (CID), with up to 20 fragment spectra (MS2) collected per MS1. Dynamic exclusion was activated, with a 45-s exclusion time for ions selected more than twice in a 30-s window.

Spectra were analyzed by SEQUEST (Proteome Discoverer 1.4, Thermo Scientific) using a protein sequence database consisting of the *E. coli* and T7 bacteriophage proteomes, including alternative extended sequences resulting from amber readthrough. These protein sequences were generated from GenBank records NC_012971.2 and NC_001604.1, respectively. The amber codon substitution in the type II holin protein was also included for phage T7. Amber sites were designated as residue X in the sequence database, with X alternatively representing 3-iodotyrosine, tyrosine, phenylalanine or tryptophan in independent searches. The search specified full-proteolytic peptides, allowing up to two missed cleavages. Mass tolerance was set to 10 p.p.m. for precursor and 0.5 Da fragment spectra. Carbamidomethylation of cysteine (iodoacetamide) was selected as a static modification, and oxidized methionine was selected as a dynamic modification. High-confidence peptide-spectrum matches (PSMs) were filtered at <1% FDR (determined by Percolator, Proteome Discoverer v1.3, Thermo Scientific). For the RF0 IodoY *E. coli*, RF0 IodoY *E. coli* cells lysed by the T7 phage from population WT L6 and BL21(DE3) *E. coli* cells lysed by wild-type T7 samples, we identified a total of 183,363, 18,527 and 20,639 PSMs, respectively. Full details of predicted PSMs that were informative about the specificity of amber readthrough are provided (**Supplementary Fig. 2** and **Supplementary Data Set 2**).

26. Dotd, M., Roehr, J., Ahmed, R. & Dieterich, C. *Biology* **1**, 895–905 (2012).

27. Gfeller, D., Michielin, O. & Zoete, V. *Nucleic Acids Res.* **41**, D327–D332 (2013).

28. Young, D.D., Jockush, S., Turro, N.J. & Schultz, P.G. *Bioorg. Med. Chem. Lett.* **21**, 7502–7504 (2011).

29. Shevchenko, A., Tomas, H., Havlis, J. & Olsen, J. *Nat. Protoc.* **1**, 2856–2860 (2006).