

# Principal Component Analysis (PCA)


BCH364C/391L Systems Biology / Bioinformatics – Spring 2015

Edward Marcotte, Univ of Texas at Austin

**What is Principal Component Analysis? What does it do?**

**So, first let's build some intuition.**

“You do not really understand something unless you can explain it to your grandmother”, Albert Einstein



wikipedia

With thanks for many of these explanations to <http://stats.stackexchange.com/questions/2691/making-sense-of-principal-component-analysis-eigenvectors-eigenvalues>

## What is Principal Component Analysis? What does it do?

### A general (and imprecise) political example:

Suppose you conduct a political poll with 30 questions, each answered by 1 (*strongly disagree*) through 5 (*strongly agree*). Your data is the answers to these questions from many people, so it's 30-dimensional, and you want to understand what the major trends are.

You run PCA and discover 90% of your variance comes from one direction, corresponding not to a single question, but to a specific weighted combination of questions. This new hybrid axis corresponds to the political left-right spectrum, *i.e.* democrat/republican spectrum.

Now, you can study that, or factor it out & look at the remaining more subtle aspects of the data.

**So, PCA is a method for discovering the major trends in data, simplifying the data to focus only on those trends, or removing those trends to focus on the remaining information.**


Example: Christian Bueno, <http://stats.stackexchange.com/questions/2691/making-sense-of-principal-component-analysis-eigenvectors-eigenvalues>

## What is Principal Component Analysis? What does it do?

### A more precise graphical example:


**In a general sense, PCA rotates your axes to “line up” better with your data.**

Because rotation is a kind of linear transformation, your new dimensions will be weighted sums of the old ones, like  $\langle 1 \rangle = 23\% \cdot [1] + 46\% \cdot [2] + 39\% \cdot [3]$



Quotes & image adapted from isomorphisms, <http://stats.stackexchange.com/questions/2691/making-sense-of-principal-component-analysis-eigenvectors-eigenvalues>

**PCA finds new variables which are linear combinations of the original variables such that in the new space, the data has fewer dimensions.**



Think of a data set consisting of points in 3D on the surface of a flat plate held up at an angle. In the original  $x, y, z$  axes you need 3 dimensions to represent the data, but with the correct linear transformation, you only need 2.

Quotes: Shlomo Argamon, from <http://stats.stackexchange.com/questions/2691/making-sense-of-principal-component-analysis-eigenvectors-eigenvalues>  
Image: <http://phdthesis-bioinformatics-maxplanckinstitute-molecularplantphys.matthias-scholz.de/>

### To summarize so far:

PCA is a technique to reduce dimension by:

1. Taking **linear combinations** of the original variables.
2. Each linear combination explains the **most variance** in the data it can.
3. Each linear combination is uncorrelated (**orthogonal**) with the others
4. Plot the data in terms of only the most important (**principal**) dimensions







Image: <http://blog.equametrics.com/2013/02/an-introduction-to-principal-component-analysis>  
Quote: <http://stats.stackexchange.com/questions/2691/making-sense-of-principal-component-analysis-eigenvectors-eigenvalues>




## Linear Transformations



**If we can stretch & rotate this way...**

Adapted from [http://www.engr.sjsu.edu/~knapp/HCIRODPR/PR\\_Maha/linear\\_tr.htm](http://www.engr.sjsu.edu/~knapp/HCIRODPR/PR_Maha/linear_tr.htm)

## Linear Transformations




**...then we can also go the other way.**

**To do this, we need to learn about covariance...**

Adapted from [http://www.engr.sjsu.edu/~knapp/HCIRODPR/PR\\_Maha/linear\\_tr.htm](http://www.engr.sjsu.edu/~knapp/HCIRODPR/PR_Maha/linear_tr.htm)

**Covariance = measures tendency to vary together (to co-vary). Similar to correlation.**



**Variance** = average of the squared deviations of a feature from its mean  
**Covariance** = average of the products of the deviations of feature values from their means

$$c(i,j) = \frac{[x(1,i) - m(i)][x(1,j) - m(j)] + \dots + [x(n,i) - m(i)][x(n,j) - m(j)]}{(n - 1)}$$

Adapted from [http://www.engr.sjsu.edu/~knapp/HCIRODPR/PR\\_Mahal/cov.htm](http://www.engr.sjsu.edu/~knapp/HCIRODPR/PR_Mahal/cov.htm)

All of the **covariances**  $c(i,j)$  between features can be collected together into a **covariance matrix C**.

This summarizes all of the correlation structure among all pairs of features.

$$C = \begin{bmatrix} c(1,1) & c(1,2) & \cdots & c(1,n) \\ c(2,1) & c(2,2) & \cdots & c(2,n) \\ \vdots & \vdots & & \vdots \\ c(n,1) & c(n,2) & \cdots & c(n,n) \end{bmatrix}$$

covariance between feature 1 and 2  
 covariance between feature 2 and n  
 etc.


## We need one last concept: Eigenvectors and eigenvalues

An **eigenvector** of a square matrix  $A$  is a non-zero vector  $v$  that, when the matrix is multiplied by  $v$ , yields a constant multiple of  $v$ , the multiplier being commonly denoted by  $\lambda$ . That is:

$$Av = \lambda v$$

(Because this equation uses **post-multiplication** by  $v$ , it describes a **right eigenvector**.)


The number  $\lambda$  is called the **eigenvalue** of  $A$  corresponding to  $v$ .



The **blue arrow** is an eigenvector of this linear transformation matrix, since it doesn't change direction.

Its scale is also unchanged, so its eigenvalue is 1.

Wikipedia




wikipedia

## Calculating the PCA

1. Calculate the covariation matrix  $C$  between features of the data
2. Calculate the eigenvectors and eigenvalues of  $C$
3. Order the eigenvectors according to the eigenvalues

PC1 is the eigenvector corresponding to the largest eigenvalue, PC2 is the eigenvector corresponding to the next largest, etc.


The data can be plotted as projections along the PCs of interest.




Here's a beautiful application of PCA that illustrates how it can simplify complicated datasets:

**What do you think are the most important genetic trends among European men?**

**What might this picture → look like?**





**Genes mirror geography within Europe**


John Novembre<sup>1,2</sup>, Toby Johnson<sup>1,3,6</sup>, Katarzyna Bryc<sup>7</sup>, Zoltán Kutalik<sup>4,6</sup>, Adam R. Boyko<sup>7</sup>, Adam Auton<sup>7</sup>, Amit Indap<sup>7</sup>, Karen S. King<sup>8</sup>, Sven Bergmann<sup>4,6</sup>, Matthew R. Nelson<sup>8</sup>, Matthew Stephens<sup>1,3</sup> & Carlos D. Bustamante<sup>7</sup>

**This is a map of their DNA. It recapitulates the map of Europe.**

**In other words, the strongest genetic trends across European men relate directly to their geographic locations (note: not nationality)**


Vol 456 | 6 November 2008 | doi:10.1038/nature07331

The trend even holds up within a country, e.g. Switzerland:




Vol 456 | 6 November 2008 | doi:10.1038/nature07331

**Why? Genetic similarity is strongly distance dependent in Europe, presumably because people tend to live near where they were born and tend to marry locally.**



Vol 456 | 6 November 2008 | doi:10.1038/nature07331

The trend is so strong that it can be used to *predict* where the men are from, e.g. shown here by leave-one-out cross-validation:




Vol 456 | 6 November 2008 | doi:10.1038/nature07331

## SUMMARY

In a sense, PCA fits a (multidimensional) ellipsoid to the data

- Described by directions and lengths of principal (semi-)axes, e.g. the axis of a cigar or egg or the plane of a pancake



- No matter how an ellipsoid is turned, the eigenvectors point in those principal directions. The eigenvalues give the lengths.
- The biggest eigenvalues correspond to the fattest directions (having the most data variance). The smallest eigenvalues correspond to the thinnest directions (least data variance).
- Ignoring the smallest directions (*i.e.*, collapsing them) loses relatively little information.

Adapted from whuber, <http://stats.stackexchange.com/questions/2691/making-sense-of-principal-component-analysis-eigenvectors-eigenvalues>



### More examples of different linear transformations:

(a) Identity  $L = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(b) Stretching  $L_1 = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$

(g) Shear and rotation  $L_6 = \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix}$

(h) Rotation and shear  $L_7 = \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}$

(c) Reflection  $L_2 = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$

(d) Stretching and reflection  $L_3 = \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix}$

(i) Rotation and stretching  $L_5 = \begin{bmatrix} 0 & -1 \\ 1 & 2 \end{bmatrix}$

(j) Stretching and rotation  $L_9 = \begin{bmatrix} -2 & 0 \\ 0 & 1 \end{bmatrix}$

(k) Projection  $L_{10} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

(l) Projection and rotation  $L_{11} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

<http://isomorphisms.tumblr.com/post/11407834056/matrices-linear-algebra>