

Synthetic Biology = design and engineering of biological systems that aren't found in nature

Why would we want to do this?

- Want to understand natural systems. One of the best ways to understand a system is to change it or make new, related ones
- To fully "understand" a system, we should be able to predict the outcome when we change the system
- For molecular biology, this means:
 - designing new gene circuits and networks
 - modeling the designed systems & predicting their properties
 - making & testing the designs
 - updating our understanding from the model/test agreement

Engineers often look at biological systems & think that the systems are equivalent to electronic circuits

e.g,

| | |
|-----------------------------|----------------------------|
| fluorescent proteins | light bulbs or LEDs |
| transcription factors | transistors or logic gates |
| repressors | NOT gates |
| activators | OR/AND gates |
| polymerases | |
| (transcriptional machinery) | batteries |



and so on...

Are they right?

→ raises the possibility that biological parts (genes, proteins, etc.) could be combined using the rules established for analog/digital circuits

The Repressilator = engineered genetic circuit designed to make bacteria glow in a oscillatory fashion = “repressor” + “oscillator”

Transcriptional repressors ↘




Elowitz & Leibler, *Nature* (2000) 403:335-8

The Repressilator = engineered genetic circuit designed to make bacteria glow in a oscillatory fashion


Repressilator

Reporter



Elowitz & Leibler, *Nature* (2000) 403:335-8

The repressilator in action...



iGEM: A synthetic biology contest

(from iGEM's web site)

Can simple biological systems be built from standard, interchangeable parts and operated in living cells? Or is biology simply too complicated to be engineered in this way?

iGEM's broader goals include:

- To enable systematic engineering of biology
- To promote open & transparent development of tools for engineering biology
- To help construct a society that can productively apply biological technology


2004: MIT, UT, Princeton, Boston University, Cornell

2005: 13 teams (the above + UK, Germany, more...)

2006: 32 teams, incl. Japan/Latin America/Korea/India/more Europe



54 teams in 2007, 84 teams in 2008, 112 teams in 2009, 130 teams in 2010, 165 teams in 2011, and 245 teams in 2012 and 2013...

UT's 2004/2005 iGEM project – build bacterial edge detector




How does edge detection work in principle?

A computer might visit each pixel in turn, and check to see if it is bordered by both black & white pixels. If yes, highlight the pixel.



Bacterial photography



Levskaya et al. *Nature*, 438(7067):441-2 (2005)



| Mask | Cph1/EnvZ |
|----------------|----------------|
| Hello World | Hello World |
| Hello World | Hello World |

"Light cannon" developed by Aaron Chevalier,
UT undergraduate

Levskaya et al. *Nature*, 438(7067):441-2 (2005)

The first bacterial photograph (coliroid?)...




Levskaya et al. *Nature*, 438(7067):441-2 (2005)

*Escherichia
darwinia*





Image: Aaron Chevalier

On to the edge detector...



It works!





UT's 2012 iGEM project – build caffeine biosensor

ACS Synthetic Biology

Letter


pubs.acs.org/synbio

Decaffeination and Measurement of Caffeine Content by Addicted *Escherichia coli* with a Refactored *N*-Demethylation Operon from *Pseudomonas putida* CBBS



Erik M. Quandt,¹ Michael J. Hammerling,¹ Ryan M. Summers,¹ Peter B. Otoupal,¹ Ben Slater,¹ Razan N. Alnahhas,¹ Aurko Dasgupta,¹ James L. Bachman,¹ Mani V. Subramanian,² and Jeffrey E. Barrick^{1,3}

Basic idea


- Block *de novo* guanine synthesis
- Convert caffeine to xanthine
- Addict *E. coli* bacteria to caffeine




ACS Synth. Biol. 2013, 2, 301–307



One major success of synthetic biology is the engineering of the Artemisinic acid production pathway from wormwood into yeast:



Increasing production of artemisinic acid by strain engineering and addition of IPM to cultures.



“These key developments in yeast strain engineering, fermentation, and artemisinin synthetic chemistry pave the way for an industrial process capable of supplementing the world supply of artemisinin from a second source independent of the uncertainties associated with botanical production.”

CJ Paddon *et al. Nature* **000**, 1-5 (2013) doi:10.1038/nature12051

Who needs nature? Made-to-order, designer organisms

Largest Gene Synthesis Supplier in USA

- 100% sequence accuracy guaranteed
- Fastest turnaround: as few as **4 business days**
- lowest price: starting at **\$0.23/bp**

www.genscript.com


**We can now manufacture a complete genome
from commodity chemicals**

**Therefore, we can program whatever changes we want,
assuming we can get it into cells...**

Creation of a Bacterial Cell Controlled by a Chemically Synthesized Genome


Daniel G. Gibson,¹ John I. Glass,¹ Carole Lartigue,¹ Vladimir N. Noskov,¹ Ray-Yuan Chuang,¹
Mikkel A. Algire,¹ Gwynedd A. Benders,² Michael G. Montague,¹ Li Ma,¹ Monzia M. Moodie,¹
Chuck Merryman,¹ Sanjay Vashee,¹ Radha Krishnakumar,¹ Nacyra Assad-Garcia,¹
Cynthia Andrews-Pfannkoch,¹ Evgeniya A. Denisova,¹ Lei Young,¹ Zhi-Qing Qi,¹
Thomas H. Segall-Shapiro,¹ Christopher H. Calvey,¹ Prashanth P. Parmar,¹ Clyde A. Hutchison III,²
Hamilton O. Smith,² J. Craig Venter^{1,2*}

“We report the design, synthesis, and assembly of the 1.08–mega–
base pair *Mycoplasma mycoides* JCVI-syn1.0 genome starting from
digitized genome sequence information and its transplantation into a
M. capricolum recipient cell to create new *M. mycoides* cells that are
controlled only by the synthetic chromosome.”




2 JULY 2010 VOL 329 SCIENCE

“Rebooting” bacteria with synthetic genomes



2 JULY 2010 VOL 329 SCIENCE

“The only DNA in the cells is the designed synthetic DNA sequence, including “watermark” sequences and other designed gene deletions and polymorphisms, and mutations acquired during the building process. The new cells have expected phenotypic properties and are capable of continuous self-replication.”



PCR of 4 engineered “watermarks”

2 JULY 2010 VOL 329 SCIENCE

But, wait! They only changed DNA, not the rest of the cell!

However...

In biology, software encodes the hardware.
Most (all?) of the cell is specified by the DNA.

It's as though you bought a Blackberry...



installed the Android operating system...

& your phone physically morphed
into a Galaxy S4...




Some good quotes from the paper:

“If the methods described here can be generalized, design, synthesis, assembly, and transplantation of synthetic chromosomes will no longer be a barrier to the progress of synthetic biology.”


“We expect that the cost of DNA synthesis will follow what has happened with DNA sequencing and continue to exponentially decrease. Lower synthesis costs combined with automation will enable broad applications for synthetic genomics.”

“As synthetic genomic applications expand, we anticipate that this work will continue to raise philosophical issues that have broad societal and ethical implications.”

In 2016, Hutchison, Chuang, *et al.* reported making living mycoplasma after cutting the genome by ½ the genes




Four design-build-test cycles produced JCVI-syn3.0. (A) The cycle for genome design, building by means of synthesis and cloning in yeast, and testing for viability by means of genome transplantation. After each cycle, gene essentiality is reevaluated by global transposon mutagenesis. (B) Comparison of JCVI-syn1.0 (outer blue circle) with JCVI-syn3.0 (inner red circle), showing the division of each into eight segments. The red bars inside the outer circle indicate regions that are retained in JCVI-syn3.0. (C) A cluster of JCVI-syn3.0 cells, showing spherical structures of varying sizes (scale bar, 200 nm).



Science 25 MARCH 2016 • VOL 351 ISSUE 6280

In parallel, methods were developed to edit genomes at many locations in parallel, e.g. reassigning all amber (TAG) stop codons in *E. coli* to ochre (TAA)



Genomically Recoded Organisms Expand Biological Functions

Marc J. Lajoie,^{1,2} Alexis J. Rovner,^{2,4} Daniel B. Goodman,^{1,5} Hans-Rudolf Aerni,^{4,6} Adrian D. Haimovich,^{3,4} Gleb Kuznetsov,¹ Jaron A. Mercer,⁷ Harris H. Wang,⁸ Peter A. Carr,⁷ Joshua A. Mosberg,^{1,2} Nadin Rohland,² Peter G. Schultz,¹⁰ Joseph M. Jacobson,^{11,12} Jesse Rinehart,^{6,8} George M. Church,^{1,3,9} Farren J. Isaacs^{1,8,4}

SCIENCE VOL 342 18 OCTOBER 2013


http://isaacs.commons.wikimedia.org/wiki/File:20131017rE.coli_Fig1.png

& now, “rebooting” yeast with synthetic chromosomes

Turns out
chromosomes can be
synthesized and
replaced for yeast too...

Synthetic Yeast 2.0
Building the world's first synthetic eukaryotic genome together

Search



Synthetic Yeast Genome, Sc2.0 2012

合成酵母基因组第一次国际会议

April 16, 2012, Beijing



& China is pushing
for a completely
synthetic yeast
genome...

Science April 4, 2014: Vol. 344 no. 6179 pp. 55-58

Total Synthesis of a Functional Designer Eukaryotic Chromosome

Narayana Annaturu,^{1*} Héloïse Muller,^{1,2,3,4*} Leslie A. Mitchell,^{2,5} Sivaprakash Ramalingam,¹ Giovanni Stracquadanio,^{2,6} Sarah M. Richardson,⁶ Jessica S. Dymond,^{2,7} Zheng Kuang,² Lisa Z. Scheifele,^{2,8} Eric M. Cooper,² Yizhi Cai,^{2,9} Karen Zeller,² Neta Agmon,^{2,5} Jeffrey S. Han,¹⁰ Michalis Hadjithomas,¹¹ Jennifer Tullman,⁶ Katrina Caravelli,^{2,12} Kimberly Cirelli,^{1,12} Zheyuan Guo,^{1,13} Viktoriya London,^{1,13} Apurva Yeluru,^{1,13} Sindurathy Murugan,⁶ Karthikeyan Kandavelou,^{1,14} Nicolas Agier,^{15,16} Gilles Fischer,^{15,16} Kun Yang,^{2,6} J. Andrew Martin,^{2,6} Murat Bilgel,¹³ Pavlo Bohutskyi,¹³ Kristin M. Boulter,¹² Brian J. Capaldo,¹³ Joy Chang,¹³ Kristie Charoen,¹³ Woo Jin Choi,¹³ Peter Deng,¹¹ James E. DiCarlo,¹³ Judy Doong,¹³ Jessilyn Dunn,¹³ Jason I. Feinberg,¹² Christopher Fernandez,¹² Charlotte E. Floria,¹² David Gladowski,¹² Pasha Hadidi,¹³ Isabel Ishizuka,¹² Javaneh Jabbari,¹² Calvin Y. L. Lau,¹³ Pablo A. Lee,¹³ Sean Li,¹³ Denise Lin,¹² Matthias E. Linder,¹² Jonathan Ling,¹³ Jaime Liu,¹³ Jonathan Liu,¹³ Mariya London,¹² Henry Ma,¹³ Jessica Mao,¹³ Jessica E. McDade,¹³ Alexandra McMillan,¹² Aaron M. Moore,¹² Won Chan Oh,¹³ Yu Ouyang,¹³ Ruchi Patel,¹³ Marina Paul,¹² Laura C. Paulsen,¹³ Judy Qiu,¹³ Alex Rhee,¹³ Matthew G. Rubashkin,¹³ Ina Y. Soh,¹² Nathaniel E. Sotuyo,¹² Venkatesh Srinivas,¹³ Allison Suarez,¹³ Andy Wong,¹³ Remus Wong,¹³ Wei Rose Xie,¹² Yijie Xu,¹³ Allen T. Yu,¹² Romain Koszul,^{3,4} Joel S. Bader,^{2,6} Jef D. Boeke,^{2,11,5†} Srinivasan Chandrasegaran^{1†}

“Here, we report the synthesis of a functional 272,871–base pair designer eukaryotic chromosome, synIII, which is based on the 316,617–base pair native *Saccharomyces cerevisiae* chromosome III. Changes to synIII include TAG/TAA stop-codon replacements, deletion of subtelomeric regions, introns, transfer RNAs, transposons, and silent mating loci as well as insertion of loxP sites to enable genome scrambling.”

Changes engineered into chromosome III

~2.5% of sequence changed

- Recoded all amber (TAG) stop codons to ochre (TAA)
- Introduced 98 Cre/Lox recombination sites
- Introduced unique sequences for PCR and new restriction enzyme sites
- Standardized telomeres

Reduced size from 316,617 bp to 272,871 bp (~14% reduction)

- Deleted 10 tRNA genes, 21 Ty elements/LTRs, silent mating loci (only one tRNA was essential, moved to a plasmid)
- Removed leucine biosynthesis gene LEU2 to be an auxotrophic marker
- Deleted all introns (affected 7 genes)
- Deleted subtelomeric DNA

Only 10 errors in assembly: 9 single base changes and 1 lost recombinase site

Last year, the Synthetic Yeast Genome Project (Sc2.0) reported on five newly constructed synthetic yeast chromosomes:



How the cover was made: <http://science.sciencemag.org/content/355/6329/eaan1126>

Design of a synthetic yeast genome

Sarah M. Richardson,^{1,2,†} Leslie A. Mitchell,^{2,3} Giovanni Stracquadanio,^{1,2,4} Kun Yang,^{1,2,†} Jessica S. Dymond,^{2,†} James E. DiCarlo,^{2,†} Dongwon Lee,^{1,5} Cheng Lai Victor Huang,² Srinivasan Chandrasegaran,² Yizhi Cai,^{2,6} Jef D. Boeke,^{2,3,‡} Joel S. Bader^{1,2,‡}

We describe complete design of a synthetic eukaryotic genome, Sc2.0, a highly modified *Saccharomyces cerevisiae* genome reduced in size by nearly 8%, with 1.1 megabases of the synthetic genome deleted, inserted, or altered. Sc2.0 chromosome design was implemented with BioStudio, an open-source framework developed for eukaryotic genome design, which coordinates design modifications from nucleotide to genome scales and enforces version control to systematically track edits. To achieve complete Sc2.0 genome synthesis,




Table 1. Design challenges and policies adopted. CDS, gene coding sequence; snoRNA, small nucleolar RNA.

| Design challenge or amendment | Policy adopted by design team |
|---|--|
| Subtelomeric repeats of varying copy number on multiple chromosomes | Delete and monitor for phenotypes as chromosomes are combined. Exception: vitamin biosynthesis genes retain one copy. |
| Dispersed repeated genes of high copy number, as well as high-copy COS and seripauperin genes | Delete and monitor for phenotypes as chromosomes are combined. |
| loxPsym sites <300 bp apart when inserted algorithmically (not especially useful and more difficult to synthesize) | loxPsym thinning to eliminate the loxPsym site closer to the centromere. |
| Stop codon overlaps a second CDS; insertion of loxPsym site would disrupt second CDS; also TAG recoding to TAA could disrupt CDS. | Favor preservation of "verified ORFs" over "dubious ORFs" and "uncharacterized ORFs"; always add loxPsym site to a verified ORF in this case |
| Tandem repeats inside CDSs (34) | Use GeneDesign's RepeatSmasher module to recode such genes to minimize DNA level repetitiveness, making DNA easier to synthesize and assemble. |
| Homopolymer tracts, including frequent A and T tracts, are difficult to synthesize | In synthesis phase, permit 10% length variation for homopolymer tracts >10 bp provided they are in a noncoding region. |
| Introns | Delete pre-mRNA introns precisely, except from genes with evidence of a fitness defect caused by intron deletion (35, 36). The <i>HAC1</i> intron, which uses separate splicing machinery and is known to play a critical role in regulation of the unfolded protein response, was not deleted (9). Delete all tRNA introns precisely. These are individually nonessential and were deleted with their host introns. |
| Intronically embedded snoRNAs | They could be "refactored" by insertion into the array of snoRNAs on chr II. |

“Mega-chunk” assembly





Table 3. Summary statistics for design of Sc2.0. WT, wild type; SYN, synthetic.

| | WT size | SYN size | No. of stop codon swaps | No. of loxP sites added | bp of PCRTag recoded | bp of RE sites recoded | No. of tRNA deleted | bp of tRNA deleted | bp of repeats deleted |
|-------|----------|----------|-------------------------|-------------------------|----------------------|------------------------|---------------------|--------------------|-----------------------|
| chr01 | 230208 | 181030 | 19 | 62 | 3535 | 210 | 4 | 372 | 3987 |
| chr02 | 813184 | 770035 | 93 | 271 | 13651 | 1215 | 13 | 993 | 7030 |
| chr03 | 316617 | 272195 | 44 | 100 | 5272 | 250 | 10 | 794 | 7358 |
| chr04 | 1531933 | 1454671 | 183 | 479 | 25398 | 2298 | 28 | 2261 | 11674 |
| chr05 | 576874 | 536024 | 61 | 174 | 8760 | 813 | 20 | 1471 | 11181 |
| chr06 | 270148 | 242745 | 30 | 69 | 4553 | 369 | 10 | 835 | 9297 |
| chr07 | 1090940 | 1028952 | 126 | 380 | 17910 | 1572 | 36 | 2887 | 13284 |
| chr08 | 562643 | 506705 | 61 | 186 | 9980 | 714 | 11 | 878 | 19019 |
| chr09 | 439885 | 405513 | 54 | 142 | 7943 | 436 | 10 | 736 | 11632 |
| chr10 | 745751 | 707459 | 85 | 249 | 12582 | 1102 | 24 | 1853 | 7523 |
| chr11 | 666816 | 659617 | 68 | 199 | 11769 | 1017 | 15 | 1243 | 4214 |
| chr12 | 1078177 | 999406 | 122 | 291 | 15129 | 1539 | 19 | 1646 | 10843 |
| chr13 | 924431 | 883749 | 100 | 337 | 15911 | 0 | 21 | 1691 | 7673 |
| chr14 | 784333 | 753096 | 96 | 260 | 13329 | 1113 | 14 | 1152 | 5115 |
| chr15 | 1091291 | 1048343 | 147 | 399 | 18015 | 2058 | 20 | 1612 | 9542 |
| chr16 | 948066 | 902994 | 127 | 334 | 15493 | 1374 | 17 | 1338 | 10048 |
| Total | 12071297 | 11352534 | 1416 | 3932 | 199230 | 16080 | 272 | 21762 | 149420 |

Synthesis, cyclization, and characterization of synV



Ze-Xiong Xie et al. *Science* 2017;355:eaaf4704

Let's end the lectures on a fun note,
with some speculative near-future
synthetic biology experiments



Science fiction? or not?
You be the judge!

“De-extincting” extinct species



Remember Dolly,
the cloned sheep?

What if the cells being cloned came
from an extinct animal and were put
into a surrogate mother?
Would that resurrect the species?

This was tried in
2009 for the
Pyrenean ibex, and
almost worked...



Cloned goat dies after attempt to bring species
back from extinction
Groundbreaking experiment fails, but scientists pave way for 'return'
of other creatures

But now there's another way!

- We can sequence a genome in a few days for a few \$K
- We can synthesize or alter big pieces of the DNA
- We can (almost) “reboot” cells with this DNA
- We can convert cells to stem cells to embryos
- We can *in vitro* fertilize animals

So why not just “edit”
the genomes of the
closest living animals to
be like their extinct
relatives?




Sound familiar?

<http://jurassicpark.wikia.com>


Besides the genome engineering, this hinges on iPS:

From embryonic stem cells, we can grow an entire organism or any cells/tissues in it



Shinya Yamanaka
Nobel Prize, 2012

& thanks to Yamanaka, we can convert skin cells back into stem cells



www.regenexx.com

There's a serious proposal to resurrect the woolly mammoth. Here's the process:

- ✓ Mammoth genome sequence
- Make ~100K DNA changes in elephant skin cells to convert elephant skin cells → mammoth skin cells
- ✓ Convert skin cells to stem cells
- ✓ Convert stem cells to embryos
- *In vitro* fertilize elephants



This might be a hard step.



www.interestingtopics.net

ANIMALS

WOOLLY MAMMOTH DNA SUCCESSFULLY SPLICED INTO ELEPHANT CELLS

BUT DON'T EXPECT MAMMOTH CLONES ANYTIME SOON

By Sarah Fecht · Posted March 24, 2015

347 Shares



Woolly Mammoth Museum

A group of researchers are

Using a DNA editing tool called CRISPR, the scientists spliced genes for the mammoths' small ears, subcutaneous fat, and hair length and color into the DNA of elephant skin cells. The tissue cultures represent the first time woolly mammoth genes have been functional since the species went extinct around 4,000 years ago.

The research has not yet been peer-reviewed or published in a scientific journal "because there is more work to do," Church told the U.K.'s *Sunday Times*, "but we plan to do so."

<http://www.popsci.com/woolly-mammoth-dna-brought-life-elephant-cells>


Which animal would you resurrect?

The dodo?



© 2010 Encyclopædia Britannica, Inc.

The quagga?




wikipedia



techandle.com

Saber-toothed tiger?

In principle, only need the DNA sequence (so, no dinosaurs)




Aurochs?

I vote for some crazy Australasian animals:

The 12'
tall
moa




& of, course, the
marsupial
Tasmanian tiger



>90° !!!

The moa-eating
Haast's eagle



What about neanderthal? Should we do it?

- ✓ Human and neanderthal genome sequence
- Edit DNA in human skin cells to convert
convert human skin cells → neanderthal skin cells
→ I give this step 10 years max before we can do this
- ✓ Convert skin cells to stem cells
- ✓ Convert stem cells to embryos
- ✓ *In vitro* fertilize
a surrogate mother

Svante
Pääbo

